

DEPARTMENT OF CLINICAL NEUROSCIENCE
Karolinska Institutet, Stockholm, Sweden

**MICROGLIA – HEALTH PROMOTING PATHWAYS AND THERAPEUTIC TARGETS IN
AGEING AND NEUROINFLAMMATION**

Rasmus Berglund



**Karolinska
Institutet**

Stockholm 2021

All previously published papers were reproduced with permission from the publisher.

Published by Karolinska Institutet.

Printed by Universitetsservice US-AB, 2021

© Rasmus Berglund, 2021

ISBN 978-91-8016-135-0

Cover illustration: By Rasmus Berglund

Microglia – Health promoting pathways and therapeutic targets in ageing and neuroinflammation

THESIS FOR DOCTORAL DEGREE (Ph.D.)

By

Rasmus Berglund

Som för avläggande av medicine doktorsexamen vid Karolinska Institutet offentligen kommer försvaras i CMM Lecture Hall, CMM L8:04, Karolinska Universitetssjukhuset Solna, 17176 Stockholm. Fredagen den 26e februari 2021 kl 09.00

Principal Supervisor:

Tomas Olsson
Karolinska Institutet
Department of clinical neuroscience
Division of Neuroimmunology

Opponent:

Manuel Friesse
University Medical Centre Hamburg-Eppendorf
Institute for Neuroimmunology and Multiple Sclerosis

Co-supervisor(s):

Maja Jagodic
Karolinska Institutet
Department of clinical neuroscience
Division of Neuroimmunology

Examination Board:

Marita Troye Blomberg
Stockholm University
Department of Molecular Biosciences

André Ortlieb Guerreiro-Cacais
Karolinska Institutet
Department of clinical neuroscience
Division of Neuroimmunology

Göran Solders
Karolinska Institutet
Department of Clinical Neuroscience
Division of Clinical Neurophysiology

Per Nilsson
Karolinska institutet
Department of Neurobiology, Care Sciences and Society
Division of Neurogeriatrics

To Moa, Edgar, Elis and Emmylou

ABSTRACT

Microglia are the innate immune cells of the CNS with an embryonic origin and self-renew with a slow turnover throughout life. The health-promoting capacities of this cell are being acknowledged through updated and specific tools separating microglia from bone marrow-derived macrophages. Knowing that microglia depends on TGF- β signaling in acquiring a mature homeostatic phenotype, we also found this cytokine to mediate the integration of monocyte-derived cells into an empty myeloid CNS niche. Microglia or the integrated monocyte-derived macrophages, absent in TGF- β signaling, developed a damaging phenotype causing spontaneous de-myelination, clinical motor deficits, and death of experimental mice.

Multiple sclerosis (MS) is a de-myelinating autoimmune CNS disease, commonly with onset in young adults as a relapsing-remitting disease that over time converts to a progressive accumulation of clinical deficits. The etiology of MS is partly inherited but, in many aspects, unknown, although we have successful treatments targeting the adaptive immune system reducing relapses and likely delay progression. However, the progressive MS disease, believed to emanate from the cells residing in the CNS, is very limited in treatment options. Genetic association studies imply that the microglial cell harness a substantial part of the MS-pathogenicity. How this relates to disease phenotypes offering treatment targets is sparsely explored. As the human CNS is rather inaccessible, the use of the rodent animal model experimental autoimmune encephalomyelitis (EAE) has been instrumental in deciphering the MS pathology. In the recovery phase of this disease, the microglial clearance of myelin is of substantial importance. We found this process to depend on a lysosomal degradation process referred to as autophagy- or LC3- associated phagocytosis. During EAE, microglia lacking the autophagy gene *Atg7* accumulated myelin debris and had reduced recirculation of scavenger receptors, causing a secondary impairment in tissue myelin-clearance. These cells also acquired an altered transcriptome associated with inflammatory microglia/macrophage phenotypes found in, e.g., MS, neurodegenerative disease, and stroke, while the genes of the homeostatic signature were downregulated. Autophagy is known to alter with age, and we targeted this by increasing autophagy-associated phagocytosis in aged microglia by treatment with the sugar molecule trehalose, which ameliorated EAE. Of note, trehalose metabolism and some autophagy-associated phagocytosis pathway components are associated with MS through risk allele analysis.

Microglial proliferation and survival rely on CSF-1 (M-CSF) or IL-34 mediated activation of the CSF-1 receptor (CSF1R). While microglial CSF-1 expression is elevated during inflammation, CSF1R associate with the homeostatic microglia. In the healthy CNS, neurons are the main source of IL-34, but a specific role of this cytokine in neuroinflammation remains to be evaluated. The ageing CNS is challenging for microglia in terms of adaptations to aid in health-promoting capacities in functions that, together with CSF1R signaling, engage canonical-autophagy. This degradation pathway controls the quality and quantity of, e.g., inflammatory mediators, receptors, and organelles. By deleting the key canonical-autophagy gene *Ulk1*, we found an age-associated subpopulation of microglia with highly activated ERK1/2 upon CSF1R engagement to be diminished, a loss not compensated by other microglia or myeloid cells. The loss of this population in aged mice caused neural and glial cell death and high mortality in EAE. In autophagy-competent aged mice, we could expand this CNS protective population specifically by IL-34 treatment and thereby ameliorate disease.

In this thesis, I present TGF- β as an essential factor in establishing a homeostatic CNS myeloid cell and a demand in aged microglia for canonical-autophagy to maintain a neuroprotective phenotype. The myelin processing by microglia through autophagy-associated phagocytosis is dissected in detail, and we can show how a decline in this pathway can be restored in aged microglia.

LIST OF SCIENTIFIC PAPERS

- I. **Berglund R**, Guerreiro-Cacais AO, Adzemovic MZ, Zeitelhofer M, Lund H, Ewing E, Ruhrmann S, Nutma E, Parsa R, Thessen-Hedreul M, Amor S, Harris RA, Olsson T, Jagodic M.
Microglial autophagy-associated phagocytosis is essential for recovery from neuroinflammation. Science Immunology, 5., 52 (2020)
- II. **Berglund R**, Guerreiro-Cacais AO, Piket E, Jagodic M, Olsson T.
The ageing CNS is protected from neuroinflammation by an autophagy dependent microglia population promoted by IL-34. Manuscript
- III. Lund H, Pieber M*, Parsa R*, Grommisch D, Ewing E, Kular L, Han J, Zhu K, Nijssen J, Hedlund E, Needhamsen M, Ruhrmann S, Guerreiro-Cacais AO, **Berglund R**, Forteza MJ, Ketelhuth DFJ, Butowsky O, Jagodic M, Zhang XM*, Harris RA*.
Fatal de-myelinating disease is induced by monocyte derived macrophages in the absence of TGF- β signaling. Nature Immunology. 19(5), 1-7 (2018).
*Equal contribution

SCIENTIFIC PAPERS NOT INCLUDED IN THE THESIS

- I. Hochmeister S, Aeinehband S, Dorris C, **Berglund R**, Haindl MT, Velikic V, Gustafsson SA, Olsson T, Piehl F, Jagodic M, Zeitelhofer M, Adzemovic MZ.
Effect of Vitamin D on Experimental Autoimmune Neuroinflammation Is Dependent on Haplotypes Comprising Naturally Occurring Allelic Variants of CIITA (Mhc2ta). **Frontiers in Neurology**. 13;11 (2020).
- II. Castelo-Branco G, Stridh P, Guerreiro-Cacais AO, Adzemovic MZ, Falcão AM, Marta M, **Berglund R**, Gillett A, Hamza KH, Lassmann H, Hermanson O, Jagodic M.
Acute treatment with valproic acid and l-thyroxine ameliorates clinical signs of experimental autoimmune encephalomyelitis and prevents brain pathology in DA rats. **Neurobiology of Disease**. 71:220-33 (2014).
- III. Guerreiro-Cacais AO, Norin U, Gyllenberg A, **Berglund R**, Beyeen A D, Rheumatoid Arthritis Consortium International (RACI), Petit-Teixeira E, Cornélis F, Saoudi A, Fournié G J, Holmdahl R, Alfredsson L, Klareskog L, Jagodic M, Olsson T, Kockum I & Padyukov L.
VAV1 regulates experimental autoimmune arthritis and is associated with anti-CCP negative rheumatoid arthritis. **Genes & Immunity** 18, 109 (2017)

CONTENTS

1. INTRODUCTION

1.1	Multiple sclerosis	1
1.2	Rodent models of multiple sclerosis	3
1.3	Microglial origin and kinetics	3
1.4	Models of microglial depletion	4
1.5	Microglial phenotypes	5
1.5.1	Homeostatic microglia and tissue interactions	5
1.5.2	TGF- β	5
1.5.3	Microglial phenotypes in CNS pathology	5
1.5.4	Phagocytosis and re-myelination in MS and EAE	6
1.5.5	The receptors regulating and executing microglial	6
1.5.6	APOE and CLEC7A	7
1.5.7	Microglia in neurogenesis and microglial neurotrophic features	7
1.5.8	Damaging and pathogenic microglial processes	8
1.6	Ageing microglia	9
1.7	Infiltrating and border-associated-myeloid cells	9
1.8	Autophagy	10
1.8.1	Canonical-autophagy	10
1.8.2	Non-canonical autophagy and phagocytosis	11
1.8.3	Autophagy in the pathology MS and other human diseases	12
1.8.4	Autophagy as a pharmaceutical target	13

2. METHODOLOGICAL CONSIDERATIONS

2.1 Mouse Cre-Lox models	14
2.2 The Diphtheria-toxin mediated depletion of microglia	14
2.3 The EAE model	15
2.4 Microglial phenotyping in vivo and in vitro	15
2.5 Autophagy monitoring and phagocytosis assays	15
2.6 Analysis of RNA sequencing data	16
3. ETHICAL CONSIDERATIONS	16
4. AIMS	16
5. RESULTS	17
5.1 Paper I	17
5.2 Paper II	19
5.3 Paper III	20
6. DISCUSSION AND FUTURE PERSPECTIVES	22
7. ACKNOWLEDGEMENTS	25
8. REFERENCES	27

ABBREVIATIONS

ALS	Amyotrophic lateral sclerosis
AMPK	5' adenosine monophosphate-activated protein kinase
ATG	Autophagy related gene/protein
BAM	Border-associated macrophages
BBB	Blood-Brain-Barrier
BMDM	Bone-marrow derived myeloid cell/macrophage
CD	Cluster of differentiation
CLEC	C-type lectin
CNS	Central nervous system
CSF	Colony stimulating factor
CX3CR1	Chemokine(C-X-C motif) 3 receptor 1
DAM	Disease-associated microglia
DCs	Dendritic cells
EAE	Experimental autoimmune encephalomyelitis
ERK	Extracellular signal-regulated kinases
IL	Interleukin
LC3	Microtubule-associated proteins 1A/1B light chain 3
MOG	Myelin Oligodendrocyte Glycoprotein
MS	Multiple sclerosis
MSR1	Macrophage Scavenger Receptor 1
mTOR	mechanistic target of rapamycin
OPC	Oligodendrocyte progenitor cell
PLP	Proteolipid protein
PPMS	Primary progressive MS
RNS	Reactive nitrogen species
ROS	Reactive oxygen species
RRMS	Relapsing-remitting MS
SLE	Systemic lupus erythematosus
SPMS	Secondary progressive MS
TAM	Tamoxifen
TGF- β	Transforming growth factor beta
T _H	T-helper cell
TNF	Tumor necrosis factor
TREM2	Triggering receptor expressed on myeloid cells 2
ULK1	Unc-51 like autophagy activating kinas


1. INTRODUCTION

1.1 MULTIPLE SCLEROSIS

Multiple sclerosis (MS) is a leading cause of neurological disability of young adults, with close to three million cases worldwide and a high and increasing prevalence in northern Europe and the US of about 1 per 750 citizens^{1,2}. In MS, an aberrant immune activation towards myelin and neuronal epitopes cause de-myelination and axonal damage leading to neurological deficits determined by the localization of the inflammatory lesion³. The variety of signs and symptoms include motor deficits, sensory losses, and cognitive impairment. They commonly appear in the age 20-40 as a Relapsing-Remitting MS (RRMS) in which a recovery phase follows clinical manifestations. In many patients, the recovery declines with age, and the symptoms get chronic as the de-myelinated lesions expand and neuronal/axonal damage accumulate, resulting in Secondary Progressive MS (SPMS). In a small proportion of patients, the recovery is absent already from the onset in a Primary Progressive MS (PPMS)³. Charcot first described MS in 1868 as a pathological entity. This was followed by a century of observations before the understanding and treatments of the disease advanced by genetic association studies, animal models, and the development of immune-modulating drugs. The more than 200 genes affecting MS susceptibility are predominantly associated with immunity, and many MS genes are highly expressed in the CNS resident microglial cell with a phenotype shared to a high degree between MS and other neurodegenerative pathologies⁴. Although the overlap of susceptibility risk genes to primary neurodegenerative diseases is small, the MS pathology includes neurodegeneration associated with oxidative stress, mitochondrial dysfunction, and elevated ion-channel activity⁵. Similar to primary neurodegenerative diseases where e.g., α -synuclein and amyloid-beta is fundamental in pathology, the protein Bassoon is recently shown to aggregate in neurons in MS, a pathology found treatable in experimental models⁶. Further, several neurotrophic pathways in myeloid-neuronal interactions, including IGF-1-and CD200R signaling, are highlighted pathways in analysis of MS risk genes⁷. In addition to the inherited genetic risk alleles, female gender, smoking, Vitamin D deficiency, obesity, and Epstein-Barr virus infection are known factors increasing MS incidence, while ageing is the strongest risk factor for disease-progression⁸.

The inflammatory MS-relapses' initiating events are likely varied, but the bouts are largely dependent on T-and B-cells specific for CNS-antigens^{3,9}. These cells evade tolerance and act synergistically in secondary lymphoid tissues to shape pathogenic T_H1 and T_H17 cells with CNS-migratory potential^{9,10}. A break-down of the blood-brain-barrier integrity is an early event in MS relapses and a prerequisite for infiltrating pathogenic peripheral immune cells, including inflammatory monocytes and adaptive immune cells. This is proposed to be an effect derived from CNS resident cells producing e.g., IL-1 β and IL-6 and direct leucocyte mediated injury³. Today, the inflammatory MS bouts are successfully treated by lymphocyte depletion through monoclonal anti-CD20 or anti-CD52 or by obstructed migration to the CNS by VLA-4 or Sphingosine-1 receptor internalization³. These treatments reduce the annual relapse-rate from approximately 0.60 to 0.15^{11,12}. However, the MS pathology involves other immune and glial cells and the MS lesions are, besides from CD8+ T-cells, dominated by macrophages of microglial or

monocyte origin¹³. The vast inflammation in MS is mirrored by elevated levels of pro-inflammatory cytokines, e.g., IFN- γ , IL-1 β , GM-CSF, IL-17, IL-23, TNF, and CXCL-13^{14,15}. Abrogating treatments or gene ablation in animal models targeting each of these cytokines ameliorate disease, but single targeting has not shown to be sufficient to treat human MS¹⁶⁻²⁷. In the progressive stages of MS, the BBB is usually more intact, and infiltration of peripheral immune cells causing relapses is rare, while other pathogenic mechanisms such as dysregulated mitochondria and iron-metabolism appear as stronger factors^{3,28}. Treatments targeting infiltrating lymphocytes have a limited effect at this disease stage, but CD20+ B-cells, plasma cells, and CD8+ T-cells remain in the CNS and reside in meninges, causing subpial cortical inflammation in interplay with parenchymal glial cells²⁹⁻³¹. MS-lesions are characterized by the activation state of macrophages and degree of demyelination, which is prominent in both acute relapse and progressive disease³². Innate immunity and target tissue interactions in general and specifically microglial phenotypes are highlighted in analyses of MS risk genes and progressive disease pathology^{7,33}. The microglial cell acts as a glial cell with neurotrophic support and as an immune cell with phenotypes associated with the initial events of MS lesions as well as progressive MS^{7,13}. This thesis aims to deconvolute and target pathological mechanisms derived from myeloid and microglial dysfunctions. Targeting these cells can potentially abate acute inflammation, but the urgent hopes lie in stimulating recovery, re-myelination, and limit axonal damage in progressive disease^{7,13,34,35}.



Multiple sclerosis

1.2 RODENT MODELS OF MULTIPLE SCLEROSIS

Since its development by Rivers et al. in the 1930s, the experimental autoimmune encephalomyelitis (EAE) models of MS have been widely used and central in understanding autoimmune de-myelinating inflammation^{36,37}. Most of the T-cell-driven pathology and pathognomonic de-myelinating processes with associated axonal damage found in human MS have been mimicked by studies in these models^{36,38}. The rodent EAE model is induced either by passive immunization in cell-transfer settings or/and by active immunization with myelin antigens or emulsified spinal cord. The myelin antigens used are complete proteins or peptides with known T- or B-cell immunoreactivity, e.g., MOG. The animal species, strains, and antigens offer a wide variety of clinical courses and reproduce specific aspects of MS pathology. In passive EAE, induction by transfer of CD4⁺ T-cells is most established, while CD8⁺ cells are abundant in lesions of both MS and some EAE models and shown to either ameliorate or aggravate disease³⁹⁻⁴². B-cells cannot transmit passive EAE, but depletion in the C57B/6 mouse MOG 1-125aa model ameliorate disease. The effect on MOG 35-55 peptide induced disease in the same strain is rather aggravating EAE, illustrating that variations in the model can reflect different aspects of human MS^{43,44}. Independently of the chosen model, the quantified clinical symptoms are usually an ascending paralysis. Besides the triggered adaptive immune response, the innate immune cells, including microglia, DCs, and monocyte-derived cells, are all fundamental in the EAE pathology, and as the focus of this thesis reviewed in depth below³³.

MS-like disease can also be induced using Theiler or murine corona viruses^{38,45}. To specifically study de- and re-myelination, the Cuprizone and the lysolecithin models are widely used where chemical damage of myelin is the initial event⁴⁶. In contrast to EAE, these models are not commonly used for the quantification of clinical motor deficits. In the studies conducted within this thesis, we employ MOG-induced EAE in C57B/6 mice.

1.3 MICROGLIAL ORIGIN AND KINETICS

The microglial cell lineage constitutes about 10% of the CNS cells and is defined by localization, function, appearance, and recently by its yolk-sac derived ectodermal origin and transcriptional profile^{34,47,48}. Microglia are commonly considered self-renewing with an annual turnover in the human CNS of about 28% with less than 1% being positive for mitotic labeling in mice and human⁴⁷⁻⁵². The establishment and maintenance of the microglia population are dependent on the transcription factors IRF8 and PU.1 and the activity of CSF1R^{53,54}. Accordingly, inactivating mutations of this receptor entail a loss of the microglial population in both humans and animal models^{49,54,55}. The CSF1R binds two known ligands; CSF-1, expressed primarily by immune cells including microglia, and IL-34, expressed by neurons and glial cells with ~ 300 times higher concentrations compared to CSF-1 during homeostatic CNS conditions⁵⁶⁻⁵⁸. IL-34 has two other known receptors expressed at low levels by microglia; Syndecan-1 (CD138) and PTP- ζ (PTPRZ1). The latter suggested to be associated with a neurotrophic phenotype, but neither have a known impact on microglial survival nor proliferation⁵⁹⁻⁶². Experimental deletion of IL-34 or CSF-1 genes reduces the population in a region-dependent manner while deleting both genes decreases microglial counts similar to the low density found in mice carrying CSF1R null mutations^{49,63-65}. The downstream effects of the CSF1R engagement are complex with the

regulatory activity of several pathways, including NF- κ B, ERK, and autophagy steering Akt and AMPK^{66,67}.

Moreover, to maintain the population, microglia are shown to undergo local stochastic expansion from differentiated cells both in health and disease, and the support for a microglial “stem-cell”, distant CNS-migration, or contribution to the population from bone-marrow-derived cells have little support⁶⁸⁻⁷⁰.

The local renewal of microglia and shaping signals from surrounding tissue gives a cue for spatially defined microglia. IL-34 is associated with the forebrain, hippocampal, and cortical microglia populations, while CSF-1 is necessary for the neurotrophic microglia population in the cerebellum and brainstem^{63,64,71}. Microglia density is higher in the IL-34 dependent regions, while microglia in CSF-1 dependent regions are more phagocytic^{72,73}. Further, neutralizing anti-CSF-1 antibodies selectively reduces the microglia population in white matter CNS, while anti-IL-34 act stronger on the gray matter population⁵⁶. These regional and possibly functionally altered microglia populations derived from IL-34 or CSF-1 is possibly attributed to variations in CSF1R affinity and binding kinetics^{63,74,75}. IL-34 and CSF-1 treatments both ameliorate EAE in a similar fashion, although a moderate but evident cytokine-specific microglial polarizing potential is reported^{57,58,75,76}.

Repopulating microglia post depletion restore the majority of the expression profile while bone marrow-derived monocytes populating the microglial niche adopt a “semi-microglia” transcriptome supporting a tissue-determined homing cue for a microglial phenotype in addition to the impact from origin⁷⁷⁻⁸⁰.

1.4 MODELS OF MICROGLIAL DEPLETION

Attempts to deplete microglia and macrophages have a long history for both scientific and therapeutic purposes. In addition to the developmental microglial loss seen in CSF1R/CSF-1/IL-34 deficient mice, microglia can also be successfully depleted using CSF1R/CSF-1/IL-34 inhibitors, clodronate liposomes, or Diphtheria toxin/-receptor or Herpes simplex receptor transgenic approaches⁸¹. If the depleting condition is released, e.g., withdrawn CSF1R inhibition or loss of induced Diphtheria toxin expression, the microglial myeloid CNS niche is rapidly reconstituted^{78,80,82}. This restored population's origin is a mixture of infiltrating bone marrow-derived myeloid cells and expansion of the microglia that evaded depletion^{78,80}. Another strategy described but rather unexplored is a transfer of cytotoxic CD8⁺ T-cells reactive to the GFP expressed in CX3CR1-GFP mice, causing an almost complete microglial depletion⁸³.

Up to date, there is no post-developmental microglia-specific depletion since the strategies presented above target bone marrow-derived macrophages and are further complicated by a “cytokine storm” and infiltration of peripheral immune cells following depletion^{84,85}. The clinical outcome upon microglial depletion in neuroinflammatory and neurodegenerative disease models is mixed which likely reflects the unsolved methodological obstacles^{76,86-92}.

1.5 MICROGLIAL PHENOTYPES

1.5.1 Homeostatic microglia and tissue interactions

In healthy conditions, the homeostatic microglia act as a glial cell with trophic neuronal support interacting with astrocytes and cells of the oligodendrocyte lineage, and in the clearance of tissue debris important in keeping inflammatory processes in check⁹³. The homeostatic microglia are characterized in mice by expression of, e.g., *P2ry12*, *Cx3cr1*, *Sall1*, and *Tmem119*^{72,94}. The human microglia is more diverse, but the homeostatic population defined by the signature expression of *P2RY12*, *TMEM119*, and *CX3CR1* is found both by single-cell RNA sequencing and mass cytometry^{72,95,96}. In addition to the homeostatic core, microglia are further transcriptionally determined by the localization and tissue conditions such as age^{72,95-97}.

1.5.2 TGF- β

The cytokine TGF- β 1 (Hereafter referred to as TGF- β) and its signaling through the TGFBR (composed of the TGFBR1 and 2 subunits) is essential in acquiring a microglial phenotype and the homeostatic transcriptome^{98,99}. The TGF- β derived signature is among CNS myeloid cells specific for microglia, and congenital ablation of *Tgf- β* entails a loss of this CNS parenchymal myeloid population¹⁰⁰. TGF- β loss does not impair cell survival in mature microglia but alters the activation to a pro-inflammatory phenotype^{98,99}. In neuroinflammation, TGF- β acts in dampening inflammation and promoting re-myelination by stimulating myelin phagocytosis in EAE^{98,101-103}. Further, TGF- β is shown to stimulate myelin phagocytosis by human macrophages in vitro¹⁰². TGFBR2 deletion causes, besides the diminished homeostatic microglial phenotype, an activating phosphorylation of transcription factor TAK1 regulating microglial derived inflammation^{99,104}.

Thus, it is evident how TGF- β has a considerable health-promoting potential to be further explored in experimental and human diseases.

1.5.3 Microglial phenotypes in CNS pathology

Upon challenges such as neuroinflammation and neurodegeneration, the genes defining the homeostatic microglia are downregulated, and an activated phenotype collectively referred to as Disease-associated microglia (DAM) is acquired⁴. These phenotypes are characterized by a changed transcriptome association with various CNS disorders and models, including MS, ALS, and other neurodegenerative conditions^{4,72,94,105}. There is no unified subset of “DAM core-genes”, but in many models, these transcriptomes are dependent on TREM2 signaling^{4,72,94,105}.

Studies of EAE and experimental neurodegeneration define DAM populations enriched for, e.g., *Ly86*, *Clec7A*, *Apoe*, *Ccl-2*, and *Lyz2* accompanied by reduced expression of homeostatic genes such as *P2ry12* and *Tmem119*^{94,105,106}. In MS, several DAM populations are identified, including a phagocytic population with an enriched expression of lysosomal component *Lamp1* and suggested to be key in re-myelination^{72,107}. Microglia found in human active MS lesions are commonly positive for macrophage markers such as p22phox, CD68, CD86, MHC class II molecules, and intracellular myelin, while anti-inflammatory markers such as CD206 and CD163 are prominent on microglia associated to inactive lesions^{13,103-108}. These markers are lost on *Tmem119*+ microglia from the chronic

lesion associated with progressive MS^{13,108,109}. Recent findings show active de-myelinated lesions microglia to stain for MERTK, TIM-3, and LAMP1, with regulatory and functional implications in phagocytosis¹⁰⁸. The DAM phenotypes associated with diseases are composed of many microglial transcriptional cues for functions that remain to be explored.

1.5.4 Phagocytosis and re-myelination in MS and EAE

Microglia are CNS resident macrophages, and phagocytosis is a key feature of both DAM and homeostatic phenotypes. Phagocytosis is the endocytic process where extracellular content is engulfed by receptor-mediated budding of the plasma membrane and degraded after fusion with a lysosome. The term macrophage describes an innate immune cell's phenotype specialized in phagocytosis but does not state their ontogeny. In MS and EAE, microglia are the paramount phagocyte of apoptotic cells and myelin debris generated by inflammation, while the damaging de-myelination is executed to a larger extent by monocyte-derived macrophages (also referred to as BMDM)^{101,110-115}. Clinical deficits in MS and EAE correlate to de-myelination of the axons, and re-myelination both protect the axons from further damage and restore the signal transduction, thus recovering function^{116,117}. The re-myelination sheets are traceable as they are shorter and thinner, but the dynamics of the opposing de- and re-myelination are challenging to study in human disease^{116,117}.

In the de-myelinating cuprizone and corona-virus MS-models and in EAE, re-myelination depends on the microglial clearance of tissue myelin debris, allowing differentiation of Oligodendrocyte-progenitor-cells (OPCs) into myelinating Oligodendrocytes (OLs)^{87,113,118,119}.

1.5.5 The receptors regulating and executing microglial phagocytosis

Most significant in dictating the phagocytic microglial phenotype in models of MS, Alzheimer's, and Parkinson's is the TREM2 receptor^{94,105,111,120-122}. In EAE, gene ablation or antibody-mediated blockade of TREM2 aggravates EAE while stimulation with an agonistic antibody increases myelin clearance and hence, OPC differentiation and re-myelination^{111,123-125}. In humans, mutations in the TREM2 gene cause microglial dysfunction and the neurodegenerative Nasu-Hakola disease^{126,127}. TREM2 recognizes phosphatidylserines. These lipids are found abundantly in myelin, and apoptotic cell membranes, and binding not only increases uptake but also stimulates the degradation of phagocytosed content^{126,128-131}. Degradation of debris and apoptotic cells is further shown to be regulated by the TAM receptors - *MerTK* enriched in the homeostatic TGF- β dependent microglia population, and *Axl* enriched in DAM^{72,94,105,132,133}. The uptake of myelin is executed by scavenger receptors MSR1 (SR-A), CR3, CD36, and possibly by TREM2 directly^{128,131,134,135}.

The myelin phagocytosis is not only an executive function of the health-promoting microglia. It is also evident how accomplished ingestion and myelin degradation further induce a beneficial microglial phenotype in MS and EAE. This phenotype is defined by transcriptome analysis and immunohistochemistry defined qualities, including secretion of TGF- β and IGF-1, a mediator central in OPC differentiation to myelinating OLs^{101,136,137}.

1.5.6 APOE and CLEC7A

Microglia exposed to apoptotic cells upregulates *ApoE* and *Clec7a* (Dectin-1) genes in transitioning from a homeostatic to a phagocytic DAM phenotype⁹⁴. APOE serves as a transporter of lipids and as a ligand for intracellular receptors such as the LXR¹³¹. This intracellular receptor and transcription factor mediate lipid efflux and initiate lipid degradation through increased expression of *ApoE* in a positive feed-back loop^{131,138,139}. The *APOE* gene contains the strongest risk allele for Alzheimer's disease and is, through binding TREM2, essential for DAM differentiation^{94,140}. The risk-variant *APOEε4* associates with dysregulated phagocytosis, exemplifying an errant DAM phenotype driving pathogenesis¹⁴⁰⁻¹⁴². Carrying the *APOE4* risk allele does not associate with MS susceptibility but is suggested to accelerate disease progression¹⁴³⁻¹⁴⁵.

As part of a microglial phenotype enriched for genes promoting phagocytosis and phagosome degradation, *Clec7a* associates with microglia important in mouse and human development^{146,147}. This microglial population is absent in the adult CNS at homeostatic conditions but shows similarities to the TREM2-dependent DAM microglia^{94,105,146,147}. The CLEC7A is commonly known to bind fungal cell wall compounds but can bind apoptotic cells during inflammatory conditions¹⁴⁸. Ablation of the *Clec7a* gene is suggested to aggravate EAE, and activation of CLEC7A by the β -glucan Zymosan promotes axonal regeneration through an ERK1/2 pathway¹⁴⁹⁻¹⁵¹.

1.5.7 Microglia in neurogenesis and microglial neurotrophic features

Microglial phagocytosis is not only a fundamental health-promoting process in the CNS during inflammation. From embryogenesis, during development, and in the adult CNS, these cells phagocytose apoptotic neural lineage cells and synapses in a process supportive of neurogenesis, myelination, and synaptic plasticity¹⁵²⁻¹⁵⁵. This process relies on the microglia population sustained by CSF1R activity and expression of the fractalkine receptor (CX3CR1), key in microglial motility and spatial orientation mediated by neuronal CX3CL1^{154,156}. The synaptic pruning in neuronal plasticity is shown dependent on activation through CR3 and TREM2^{157,158}. On the negative side, in the aged CNS, excessive synaptic clearance by complement receptor-activated microglia associate with cognitive decline^{159,160}. Neuronal-microglial interaction through CD200-CD200R stabilize microglia and possibly act trophic on neurons¹⁶¹. Dysregulation in this and the CX3CL1-CX3CR1 communication is found in MS lesions, Alzheimer's disease and in the ageing brain^{161,162}. Disruption of the CX3CR1 signaling also attenuates experimental neuroinflammation, although this effect is not defined as specifically microglia derived^{163,164}.

Pathway analysis of the risk-genes associated with MS highlights the IGF-1, CNTF, BDNF, EGF, IL-10, and NGF neurotrophic pathways, all mediators secreted by microglial cells during inflammation⁷. Of certain interest is *Igf1* found enriched in DAM, especially in the phagocytic microglia population associated with EAE recovery, the anti-inflammatory cytokine IL-10 secreted upon phagocytosis, and BDNF with a known age-associated decline^{101,105,165-167}. Microglia is in addition to this neurotrophic support shown to recruit neuronal progenitors in the adult CNS in response to tissue damage¹⁶⁸.

1.5.8 Damaging and pathogenic microglial processes

Microglia exhibits damaging potential through cytokine and ROS/RNS secretion during inflammation or neurodegeneration, and microglial depletion can ameliorate pathology^{90,91}. During neuroinflammation, ROS is generated in myeloid cells by myeloperoxidases, NOX-complexes, and mitochondria^{169,170}. Errant ROS production is an early and pathogenic event in autoimmune neuroinflammation and is counteracted by redox mechanisms, e.g., NRF2 mediated expression of antioxidants^{13,170-172}. Moreover, catalase treatment reduces ROS and stabilizing the mitochondria, protects the axons, and ameliorates EAE¹⁷³⁻¹⁷⁶. Extracellular ROS is damaging myelin and axons and acts on the BBB to increase permeability^{177,178}. Abrogated ROS production through NADPH inhibition reduced infiltration of peripheral immune cells and ameliorated EAE^{177,178}. In contrast, intracellular ROS production from CNS macrophages is induced by phagocytosis, and phagocytic clearance of myelin is suggested to demand ROS^{179,180}. The effects of oxidative stress are, however, complex, and the MS drug dimethyl fumarate is shown to, besides its NRF2 activating abilities, also to increase peripheral oxidative stress in an immunomodulatory effect¹⁸¹.

Re-myelination is regulated by myelin debris-clearance but also by cytokines secreted by glial cells and immune cells. The differentiation of OPCs and thus re-myelination is inhibited by TNF, IL-17, and IFN- γ found in MS lesions and produced by microglia, although they are not considered the primary source¹⁸²⁻¹⁸⁴. Macrophage NLPR3 inflammasomes are found abundant in lesions of RRMS and primary progressive MS where IL-1 β is a candidate biomarker for poor prognosis and possibly a treatment target¹⁸⁵. In EAE, IL-1 β has an ambiguous role, damaging through recruiting and polarizing T-helper cell phenotypes driving neuroinflammation. However, IL-1 β also attracts and stimulates differentiation of OPCs and thus exhibits a positive impact on re-myelination¹⁸⁶⁻¹⁸⁹. This is partly mediated through the induction of microglial secretion of LIF and IGF-1 acting on OPCs^{188,189}. However, this is a complex issue, and IL-1 β is also shown to delay re-myelination in the Cuprizone de-myelination model¹⁸⁵.

Microglia are found early in MS lesions and participate in the recruitment and polarization of pathogenic peripheral immune cells by being a cytokine-source of, e.g., CXCL13 recruiting B-cells to the CNS, IL-1 β and IL-23 polarizing pathogenic T-cells and allowing for monocytes to enter the CNS^{17,23,24,186}. All with a crucial impact on EAE and correlating to MS disease^{14,16,17,19,24,190}. The TAK1-Nf κ B axis regulates microglial cytokine expression that is a prerequisite for immune cell infiltration and de-myelination in developing EAE¹⁰⁴. These events are at the crossroads of the “inside-out” vs. “outside-in” theories, where the latter represent the established idea that the event initiating a relapse is peripheral immune cells evading tolerance and entering the CNS¹⁹¹. The alternative “inside-out” theory claims that myelin damage is the initial event followed by microglial activation and recruitment of peripheral immune cells, accelerating the damage¹⁹². Support for this comes from animal studies with induced oligodendrocyte or myelin damage leading to inflammatory relapses and rare mutations in a myelin component gene associating with MS^{192,193}.

1.6 Ageing microglia

The Ageing of an individual is not a uniform process in all its organs and cells. Microglia are cells with low mitotic activity, which keep the telomeric length but, on the other hand, opt out on the possibility of a fresh start as the cells accumulate misfolded proteins and leaky mitochondria¹⁹⁴. Age is associated with an altered immune system, dysfunctional myeloid cells, neurodegeneration, and for MS patients associates with a switch to progression¹⁹⁴⁻¹⁹⁶. The core homeostatic microglial phenotype of adult mice in healthy conditions is also well represented in aged mice and humans, and the transcriptomes overlap between the species^{97,197,198}. However, in both mice and humans, advanced age is associated with alterations in microglial subpopulations where transcriptomes indicate reduced signaling of the TGF- β pathway and increased expression of genes involved in vesicle biogenesis and phagocytosis, including *APOE*^{197,198}. Despite this, functional microglial phagocytosis of protein aggregates and myelin debris decline with age while the lysosomal burden increases indicating either a dysfunctional degradation or higher degradational demand¹⁹⁹⁻²⁰¹. The aged microglia accumulate leaky ROS-producing mitochondria and display NLRP3 inflammasome instability, both pathogenic in progressive MS and neurodegeneration^{185,202,203}. The age-associated changes in lysosomal degradation and ROS production associate with a so-called Lipofuscin accumulation of highly oxidized protein aggregates that in microglia cause further defects in metabolism, inflammatory polarization, altered morphology, and reduced survival^{77,200,204,205}. Of note, these phenotypical changes and the microglial senescence could be derived from an age-associated autophagy impairment, which is discussed in depth later.

As a cure for an age-associated decline in microglial function, an aged microglia replacement is proposed. In experimental models, the repopulating microglia had reduced cellular and tissue ageing features, including a lowered load of lipids and lysosomes and reduced *ApoE* expression^{77,206-208}. The re-established microglia is further shown to differentiate accordingly to CNS-region defined cues^{208,209}. The clinical impact from a true microglia replenishment of the myeloid CNS niche in EAE and MS remains to be evaluated.

1.7 Infiltrating and border-associated-myeloid cells

Microglia are the only CNS parenchymal myeloid cells in homeostatic conditions, but macrophages of other origins populate the CNS interfaces¹⁰⁶. These populations, collectively referred to as BAMs or CAMs (Border or CNS Associated Macrophages, respectively; hereafter referred to as BAMs), reside in the perivascular space, the choroid plexus, and in the meninges. By fate-mapping, these cells are suggested to derive from an embryonic CD206+ population giving rise to BAM, while the CD206- is the origin of the TGF- β dependent microglia population²¹⁰. Similar to microglia, BAMs rely on the transcription factors PU.1 and IRF8. These cells are self-renewing in homeostatic conditions, while depletion through CSF1R-blocking resulted in persistent contamination with bone marrow-derived myeloid cells^{210,211}.

The BAM populations all have transcriptomes disparate from microglia except a small subset of macrophages residing in the choroid plexus^{210,211}. This population shares the transcriptional signature with TREM2 derived DAM, suggesting the existence of a non-parenchymal microglia-like cell, possibly the cell known as “Kolmer’s epiplexus cell”

described first in 1921^{211,212}. As with other DAM's, these cells have enriched expression of genes involved in phagocytosis and lipid metabolism, including *Clec7a* and *ApoE*, although in homeostatic conditions²¹¹. The BAMs are all targeted in the CX3CR1 inducible CRE model, but compared to microglia, they constitute tiny populations with no solid evidence of impact on MS and EAE, although they respond to experimental neuroinflammation by altered expression patterns^{106,213}. However, the meninges are considered a niche for lymphoid cells in progressive MS, and the choroid plexus is suggested to be a route of entry of T-cells in MS, implying these populations to influence a disease pathology not fully covered by animal models^{214,215}.

The inflamed CNS tissue in MS and EAE is infiltrated by classical and nonclassical monocytes differentiating into monocyte-derived dendritic cells and macrophages. Monocyte-derived DCs have an EAE-specific signature with, e.g., elevated expression of *Clec9a* coding a receptor binding necrotic cells for processing and cross-presentation^{106,216}. The monocyte-derived macrophages are distinguished from microglia by, e.g., surface marker CD49d or high expression of CD44, CD45, or F4/80 and are functionally less health-promoting phagocytes during EAE^{18,78,106,110,114,217}. Although microglia have pathogenic potential, executive functions in T-cell activation, de-myelination, and axonal damage largely depend on CCR2+ and/or Ly6C+ infiltrating bone marrow-derived myeloid cells^{18,114,186,218-220}. In EAE, these cells egress from bone-marrow and enter the CNS in a CCL-2 and GM-CSF dependent manner^{18,21,186,219-222}. The pluripotent cytokine GM-CSF is found abundantly in MS lesions and demanded in EAE induction, and overexpression causes spontaneous neuroinflammation with infiltration of inflammatory monocytes^{20,21,186,221,222}. Of note, these infiltrating myeloid cells do not seem to contribute to the CNS parenchymal myeloid population once the autoinflammatory EAE relapse is ceased²²⁰. The “M1” and “M2” macrophage nomenclature has not survived the complexity added by methodological advances but assigned “M2” bone marrow-derived macrophages such as CD206+ ARG1+ IGF1+ cells are associated to re-myelination, and transfer of TGF-β stimulated macrophages ameliorates EAE^{119,223}.

Genes involved in antigen presentation, including the pivotal risk allele HLA DRB1*1501, are the strongest risk factors for MS where T-cells activated by APCs are pathognomonic³. Microglia are known to be weak in APC functions in stimulating and polarizing T-cells, a function rather executed in the CNS in MS and EAE by monocyte-derived DCs and macrophages^{106,224-227}. Before entering the CNS, T-cells are activated in lymphoid tissues and/or possibly by the BAMs or DCs in the choroid plexus^{214,226,228}.

We can conclude that the myeloid CNS landscape can be far more complex and adaptive than previous notions, with functions defining subpopulations and associations to pathology to be discovered.

1.8 AUTOPHAGY

1.8.1 Canonical autophagy

Canonical (macro)-autophagy is by definition a catabolic process degrading errant or superfluous intracellular proteins or organelles in an endosomal-lysosomal route. This process uses a set of proteins, commonly prefixed ATG, with defined functions at the different autophagosome formation steps. While the energy state of a cell and its surrounding tissue is the foremost regulator of autophagy, many other intra- and extra-

cellular stimuli, including CSF1R, TREM2, and C-type lectin receptor engagement, also regulate myeloid cell autophagy^{75,229-231}. The launching step of autophagosome formation is the pre-initiation complex assembly, a process induced by AMPK kinase activity and regulated negatively by Akt by activating the mTOR-complex. These pathways are regulated downstream the CSF1R and TREM2 and induce autophagosome maturation in myeloid cells by activating ERK1/2^{66,128,130,232,233}. TREM2 is also shown to inhibit aberrant mTOR-mediated autophagy in a model addressing this receptor as a risk factor for Alzheimer's disease²²⁹. The aged microglia is shown to depend on alterations in mTOR-signaling in acquiring the age-associated phenotype both in human and animal models²³⁴.

In canonical autophagy, the ULK1/2 acts as a regulated and regulating node acting downstream by phosphorylating core-autophagy proteins, including activation of the BECLIN1/VPS34 nucleation complex leading to formation the phagophore membrane²³⁵. With the vesicle membrane in place, the cargo for degradation is loaded in either a selective or non-selective process. The phagophore maturation to a double-layer autophagosome requires ATG7 activation of the ATG5-ATG12 complex and a cysteine residue's lipidation on MAP1LC3 (commonly referred to as LC3-II when lipidated) exposed by the ATG4 kinase. LC3 is moreover suggested to have an additional role in the selection of cargo²³⁶. The LC3-II coat the autophagosome and is used as the standard defining marker of autophagy. A RAB GTPase and SNARE protein mediate the fusion with a lysosome that finalizes the degradation of the autophagosome's and its cargo²³⁷. By being the main route of degrading various cellular components, autophagy is instrumental in cellular plasticity and survival, emphasized by the findings that mice carrying *Ulk1/2*, *Atg5*, or *Atg7* constitutive gene deletions are neonatally lethal²³⁸.

Autophagy is generally considered to promote survival and differentiation in immune cells, while inhibition is associated with proliferation, including malignant myeloproliferation^{239,240}. As many of the immune system mediators regulate autophagy, autophagy also controls immune responses by regulating the density of immune receptors and providing peptides for antigen-presentation exemplified by a model where *Atg5* deficiency specifically in dendritic cells ameliorates EAE^{228,241,242}. Autophagy directly regulates inflammasome stability and secretion of IL-1 superfamily cytokines, where IL-1 β is well known in MS and EAE. Further, impaired canonical autophagy is entailed to harmful inflammatory processes, including ER stress and defective clearance of mitochondria causing dysregulated ROS production and increased Lipofuscin load^{204,243-247}. Lipofuscin associates with cellular senescence which impairs function and reduces microglial density in chronic MS lesions^{13,248}.

1.8.2 Non-canonical autophagy and phagocytosis

Non-canonical autophagy is a loose definition of cellular vesicular formations using parts of the autophagy protein machinery. This nomenclature includes activities not traditionally addressed as autophagy, such as microbe degradation in "Xenophagy" and ATG dependent secretion of cytokines. In a process referred to as autophagy- or LC3-associated phagocytosis, myeloid cells load dying cells, debris, or microbes into phagosomes labeled by LC3-II for degradation through the autophagosomal-lysosomal path²⁴⁹. This process is induced by binding to a selection of myeloid surface receptors, including CD11b, TLR4, and phosphatidylserine receptors, including Tim 3/4 and Fc gamma receptor - all myeloid


receptors implicated in neuroinflammation²⁵⁰. Further, the engagement of CLEC7A is known to induce LC3-associated phagocytosis during fungal infections^{251,252}. Autophagy-associated phagocytosis is orchestrated by the protein RUBICON that simultaneously inhibits canonical autophagy by restraining VPS34²⁵³. This process is not regulated by mTOR and independent of the pre-initiation complex containing ULK1/2 but relies on the enzymatic activity of ATG7 in lipidation of LC3^{249,253,254}. With the autophagy-defining lipoprotein embedded in the phagosome membrane, the lysosome fusion is facilitated, and phagocytosed content degraded^{249,253,254}. In addition to LC3-II, degradation of this single-layer auto-phagosome require ROS production from the NOX2 complex^{228,253}. Moreover, halted degradation of these vesicles also influences clearance capacity by impeding retromer trafficking of scavenger and regulatory receptors, e.g., CD36 and TREM2^{134,255,256}.

Impairment of the macrophage autophagy-associated phagocytosis is shown to aggravate inflammation and autoimmunity in an antibody mediated lupus-like disease and in models of autism, Alzheimer's, and Parkinson's disease with phenotypes derived from microglia deficiency in *Atg5* or *Atg7*^{255,257-259}. In these models, pathology associates both with errant tissue clearance of proteins and debris and with other functional alterations of the targeted microglia and macrophages, e.g. cytokine secretion^{253,255,257,260}.

1.8.3 Autophagy in the pathology MS and other human diseases

Genetic alterations and mutations causing dysregulated or functional impairment of autophagy are found in several human pathologies ranging from asthma to malignancies. Among these diseases, we find several neurodegenerative conditions such as ALS, Parkinson's, and Alzheimer's disease, where many risk genes associated with impaired degradation of mitochondria and errant ROS production²⁶¹. The accumulation of neuronal protein aggregates in neurodegenerative diseases such as Alzheimer's, ALS, and Parkinson's is suggested to be a result of dysfunctional autophagy-associated clearance²⁶²⁻²⁶⁴. In Crohn's inflammatory bowel disease, an *ATG16L1* risk allele associates with increased mucosal inflammation^{261,265}. This autoimmune disease entails an increased risk for developing MS and shows overlapping pathology to the one seen in the de-myelinated CNS^{255,257}. Further, an impairment specifically in autophagy-associated phagocytosis is suggested in SLE pathology where an *ATG5* allele increases risk supposedly through errant clearance and processing in phagocytosis of dying cells^{257,266}.

In MS *ATG4D*, involved in LC3 labeling of autophagosomes in all forms of mammal autophagy, is detected as a risk gene for susceptibility⁷. Also, phagosome formation and maturation are detected as regulated pathways and *RAB5*, a GTPase important in phagosome maturation in, e.g., LC3 associated phagocytosis, have a highly significant risk-allele for MS incidence^{7,253,255}. Autophagy regulatory mTOR, AMPK, Akt, and ERK pathways are also detected in pathway analysis of MS risk-genes⁷. Moreover, *CLEC16A* risk alleles in MS susceptibility associate mechanistically to MHC class II vesicular compartment maturation in a process suggested being autophagy dependent^{261,267-270}. Finally, histopathological examination of microglia in MS lesions shows the increased density of autophagosomes and NOX2 activity, possibly linked to autophagy-associated phagocytosis^{109,127}.



Autophagy

1.8.4 Autophagy as a pharmaceutical target

Autophagy is a major house-keeping pathway in most mammal cells, and pharmaceutical regulation is applicable both in, e.g., stimulated degradation of hazardous proteins, inhibiting tumor cell survival processes, and modulating the immune response, including IL-1 β and ROS secretion. However, targeting of an essential function such as autophagy could potentially increase the incidence of adverse effects²⁷¹. The well-known autophagy-inducer rapamycin reverses the mTOR suppression on autophagy and acts as an immune suppressant preventing allograft rejection. Rapamycin also ameliorates EAE partly through ERK1/2 activation and is recently shown to alter the age-associated microglial phenotype^{272,273}. Spermidine, found in, e.g., cheese, grains, and peppers, induces autophagy through downstream deacetylation of several ATGs and has been shown to increase lifespan in flies, worms, mice, and possibly humans mimicking benefits of calorie restriction^{274,275}. In the MS-model EAE, spermidine alleviates clinical symptoms and pathology by polarizing monocytes to ARG1⁺ macrophages with reduced IL-1 β secretion^{276,277}.

Another autophagy regulation goes through the activation of TFEB, a transcription factor stimulating autophagosome and lysosome component formation providing substrates for both canonical and non-canonical pathways shown protective in neurodegeneration and atherosclerosis through microglia and macrophage functions²⁷⁸⁻²⁸¹. In these studies, TFEB activation is obtained upon treatment with trehalose, a sugar-molecules naturally occurring in, e.g., shellfish and mushrooms^{279,282}. Trehalose is shown to induce autophagy through blocking surface glucose transport receptors on hepatocytes but how this associate with


TFEB and its relevance in myeloid cells remains to be explored²⁸³. Like spermidine, trehalose is not only suggested to have therapeutic potential for specific pathologies but also to act in general for CNS health and long life^{274,282,284}. Importantly, trehalose metabolism is found associated with MS through a risk allele of the gene coding for the trehalose degrading enzyme trehalase^{7,285}.

The genetic MS-associations indeed put the autophagy pathway and potential influence of environmental factors in the form of trehalose in an excellent position for further work, perhaps also therapeutically.

2 METHODOLOGICAL CONSIDERATIONS

2.1 Mouse Cre-Lox models

All experiments in **Paper I, II, III** were conducted using C57BL/6 mice. Gene ablation was accomplished using mice with cell-specific expression of the CRE-enzyme excising targeted genes flanked by lox(P) sites, referred to as Flox/FI/fl. In homeostatic conditions, Cre expressed under the *Lyz2* promoter targets primarily bone marrow-derived myeloid cells, and surprisingly also neurons²⁸⁶. Microglial *Lyz2* expression and Cre-derived deletion are increased during inflammation¹⁰⁶. Tamoxifen-induced *Cx3cr1^{CreERT2}* (hereafter referred to as *Cx3cr1^{CreER}*) model has a high recombination frequency in the microglial population but is only specific when bone marrow-derived populations are replaced from the periphery, which is not necessarily the case for BAMs^{210,287,288}. Considering this and the possible impact on phenotype from Tamoxifen itself, experiments were conducted at least 4 weeks after the last Tamoxifen treatment. In this strain, CX3CR1^{CRE} is fused with a yellow fluorescent reporter (eYFP) reporter.



*Microglial targeting in the *Cx3cr1^{CreER}* model*

2.2 Diphtheria-toxin mediated depletion of microglia

Paper III is an application of thoroughly explored possibilities of depleting the microglial population⁷⁸. The most successful depletion was accomplished by *Cx3cr1^{CreER}* mediated deletion of a stop cassette in Rosa26^{DTA} mice, leading to cell-specific diphtheria toxin

expression, which induces apoptosis and depletion of >99% of CNS myeloid cells. This could be maintained by Tamoxifen treatment-induced Cre expression. By transferring *Lyz2^{Cre}Tgfb²^{fl/fl}* or *Cx3cr1^{CreER}Tgfb²^{fl/fl}* to irradiated *Cx3cr1^{CreER}R26^{DTA}* recipient mice creating a chimera, a depletion was combined with a specific gene deletion in the monocyte and BMDM populations.

2.3 The EAE model

The EAE model employed in these studies was induced by active immunization with mouse recombinant MOG 1-125aa protein. The de-myelination occurs primarily in the spinal cord upon infiltration of peripheral immune cells and activation of microglia. This pathology includes inflammation, de-myelination and axonal damage/neuronal death. The induction of active EAE in C57BL/6 requires Freund's complete adjuvant containing *Mycobacterium tuberculosis* which leads to pattern recognition receptor activation and amplification of the immune response. Immunization is accompanied by intraperitoneal injections of *Bordetella pertussis* toxin, which induces myeloid cell IL-1 β secretion and possibly regulate BBB permeability^{19,289-291}. In **Paper I** and **III**, mice of advanced age were used. Aged mice are known to have heterogeneous phenotypes and develop aggravated EAE upon MOG immunization.

2.4 Microglial phenotyping in vivo and in vitro

In our studies, dynamics of microglial phenotypes were evaluated by transcriptional analysis, ELISA cytokine quantification, flow cytometry, functional assays, and immunocytochemistry. Disease processes were examined and visualized by immunohistochemistry of CNS with microglia and macrophages at the site of pathology. Tools for studying specifically microglia and exclude bone marrow-derived macrophages are still developing. While microglia in flow cytometry can be defined as CD45^{Intermediate} CD11b⁺ Ly6G⁻ under homeostasis, in the inflamed CNS no definite microglial marker for immunohistochemistry analysis is available. Healthy conditions offer, however, several microglia-specific markers such as TMEM119, SALL1 and P2RY12. Of note, the microglia signature is rapidly lost when cells are extracted from the CNS and cultured in vitro as they acquire an activated age-associated phenotype²⁹². Time in culture and in vitro experiments were due to this kept to a minimum.

2.5 Autophagy monitoring and phagocytosis assays

In **Paper I** and **II**, autophagy was targeted by deletion of core autophagy genes *Ulk1* and *Atg7*. *Ulk1* has high homology with *Ulk2* also expressed in microglia, but the up-, and down-stream regulation differs between the paralogs^{60,61,293}. The consequence on autophagocytosis from these deletions was assessed as, e.g., IL-1 β secretion, mitochondrial load, and LC3-II⁺ autophagosome densities after starvation phagocytic intake. Myelin-containing phagosomes targeted for degradation through the autophagy-lysosomal pathway were detected by membrane-bound LC3-II in an analysis where a mild permeabilization protocol allowed for unbound LC3-I to diffuse from the cytosol. In **Paper I**, we designed an experiment for flow cytometry to analyze the intracellular localization of phagocytosed content. Myelin or apoptotic cells were conjugated to one dye fluorescent in low pH

(lysosomes) and one emitting in a pH-independent manner (vesicles or cytosol). These experiments and most phagocytosis quantification detect uptake as intracellular content such as myelin or bacterial compounds, which is misleading regarding phagocytic clearance if the degradation of phagosomes is errant. To address this, we measured the capacity to clear myelin debris by exposing microglia and bone marrow-derived macrophages to fluorescent myelin, followed by quantification of the remaining myelin in the culture medium.

2.6 Analysis of RNA sequencing data

In **Paper I** and **III**, comprehensive expression analyses of RNA sequencing data were used to phenotype microglia. Differentially expressed genes (DEGs) were analyzed and visualized by detection of regulated pathways and pathologies in Ingenuity pathway (IPA) analysis, REViGO, and Over-representation analysis (ORA). We further employed a Gene Set Enrichment Analysis (GSEA) to investigate differences in expression between strains regarding predefined expression signatures. This was employed to compare our sequencing data to DAM-subpopulation signatures defined by single-cell RNA sequencing of microglia in, e.g., MS and Alzheimer's disease models. However, our experiments aimed not to identify new subpopulations that arise in the models deficient in targeted genes but rather to contextualize the general shifts in microglial phenotypes.

3. ETHICAL CONSIDERATIONS

For obvious reasons, CNS tissue from humans is not easy to access, and animal experiments have been instrumental in finding cues in treating and understanding neurological disease and treatment targets. Experimental animal research entails a responsibility to ensure quality and careful planning to reduce suffering. I have, through the Ph.D. education kept the use of animals to the minimum without hampering statistical significance. During the later parts of my education, I refined my technical skills and could use the same animal for several experimental analyses. To replace in vivo and ex vivo experiments was not possible due to the tissue-dependent nature of targeted microglial cells. All experiments have been conducted according to the ethical permits.

4. AIMS

Paper I

To characterize how impaired autophagy by gene ablation or age-associated decline influence cellular phenotype and clinical EAE. We further aimed to explore if this pathway could be targeted pharmacologically to ameliorate disease.

Paper II

To investigate if microglia depend on autophagy in maintaining a CNS protective population during ageing.

Paper III

To study if peripheral cells depend on TGF- β in the process of integrating to the microglial CNS niche and phenotype.

5. RESULTS

5.1. Paper I

Previous studies by our group associated reduced *Atg7* expression in immune cells to aggravated EAE in an unbiased approach in rat-strain crosses²⁹⁴. We induced *Atg7* deficiency in T-cells under the *Lck* promoter and myeloid cells under the *Lyz2* promoter in a mouse Cre-Lox system. The deletion in T-cells had minor effects on CD8⁺/CD4⁺ T-cell proportions in lymph nodes and did not affect clinical EAE. The *Lyz2*-CRE driven *Atg7* deletion was associated with a loss of recovery from EAE. Since *Lyz2* is expressed in many of the myeloid cells involved in EAE, we quantified *Atg7* expression in different subsets and found *Atg7* highly expressed and regulated in the CNS resident myeloid cell microglia. We also found deletion of floxed *Atg7* in microglia under the *Lyz2* promoter to be elevated during inflammation. To further explore this, we generated a model with microglia-specific deletion of floxed *Atg7* using the *Cx3cr1^{CreERT2/+}* (hereafter referred to as *Atg7^{fl/fl}Cx3cr1^{CreER}*), which have the benefit of an inducible deletion. This is important since autophagy has many key functions in the development, differentiation, and survival of the cell, tissue, and individual.


The clinical EAE in the *Atg7^{fl/fl}Cx3cr1^{CreER}* reproduced the chronic course seen in the *Lyz2^{CRE}* model, supporting a notion of a microglia-derived phenotype in both strains. In MS and EAE, phagocytic clearance of dead cells and myelin debris is suggested to be a microglial duty, and microglia lacking *Atg7* had accumulations of intracellular myelin. Compared to Wt, we found the myelin-containing phagosomes in autophagy-impaired microglia to have reduced LC3-II labeling and impaired loading into low-pH lysosomes. This concludes a dysfunctional phagocytic degradation through LC3-/autophagy-associated phagocytosis, a process previously known in macrophage context in models of SLE and Alzheimer's disease^{255,257}. This process was discriminated from canonical autophagy by employing deletion of *Ulk1*, an autophagy protein not demanded in autophagy-associated phagocytosis. Further, the expression of the master regulator of autophagy-associated phagocytosis *Rubicon* was found upregulated early during EAE, which supported the importance of this process. In a myelin clearance assay, we aimed to explore if the impaired degradation affected the uptake and hence clearance of tissue debris. The *Atg7*-deficient microglia had a rapidly exhausted capacity of myelin clearance, and after exposure to myelin debris in vivo and in vitro, they accumulate not only the content but also myelin scavenger receptors MSR1 (SCARA1, CD204, SR-A) and CD36. We further validated this path by detecting MSR1 blockade to inhibit myelin uptake and blocked lysosomal fusion to reduce receptor-recirculation.

Analyses of EAE CNS detected an increased intracellular myelin load in microglia lacking *Atg7* accompanied by an increased load of tissue myelin debris and consequential reduction in OPC differentiation to myelinating cells. During EAE, the *Atg7^{fl/fl}Cx3cr1^{CreER}* mice had increased immune cell infiltration to the CNS, including bone marrow-derived macrophages. However, these cells could not compensate for the reduced phagocytic capacity of microglia, supporting the idea of microglia as the primary myelin phagocyte during CNS inflammation. RNA sequencing of microglia sampled during EAE associated a DAM signature to the *Atg7* deficiency. Accordingly, we found upstream DAM regulators TREM2 and *Apoe* highly expressed in *Atg7^{fl/fl}Cx3cr1^{CreER}* microglia during EAE. We could also define and validate true subpopulations based on CLEC7A, one of the DAM signature

genes. The CLEC7A^{High} population was increased in counts in the CNS of *Atg7^{fl/fl}Cx3cr1^{CreER}* mice and possessed this model's highlighted impairments, including accumulated intracellular MSR1. Pathway analysis of microglia during EAE revealed activated inflammatory pathways detected as enriched IFN γ and GM-CSF signaling in *Atg7* deficient microglia at day 21 p.i. The later day 35 p.i. timepoint showed regulation of, e.g., metabolism and oxidative stress pathways associated with progressive MS^{3,5,115}.

Autophagy is a process known to decline with age, which also is a strong risk factor for progressive MS. As discussed above, several genes implicated in autophagy and phagocytosis have allele variants associating with MS risk⁷. By treating aged mice with trehalose, a sugar molecule found in, e.g., plants, fungi, and shellfish, we could reverse the aggravated EAE associated with age in a microglia-dependent fashion. Trehalose induced TFEB nuclear transfer and thus transcriptional activity, increasing the biogenesis of autophagosomes and lysosomes. Our experiments found a specific effect on autophagy-associated phagocytosis as the effect was not abrogated by ULK1 deficiency.

This study shows how microglia regulates EAE recovery through phagocytosis of myelin and associated phenotypic alterations. We also showed how age-associated impairments in this process could be reversed by a dietary sugar compound potentially affecting a pathway untargeted by available medications and possibly of certain importance in age-associated progressive MS.



Graphical abstract for Paper I

5.2 Paper II

As age entails altered canonical-autophagy and microglial phenotypes, we sought to examine this in the context of CSF1R signaling and neuroinflammation^{4,234,295,296}. In the aged CNS (>20 months), we found microglia to have a reduced proliferation tied to an increased cell survival indicating a slower turnover compared to adults (<5 months). While CSF1R, the main receptor for microglial survival and proliferation, had reduced surface density on aged microglia, the corresponding ligands CSF-1 and IL-34 had elevated expression in the aged CNS. By analyzing known phosphorylation-sites downstream of CSF1R engagement, we found aged microglia to have increased activating phosphorylation of ERK1/2, AMPK, and a reduced activating Akt phosphorylation. Consequently, aged microglia were significantly more sensitive to ERK1/2 inhibition. AMPK induces canonical-autophagy through the mTOR and pre-initiation autophagosome complex, including ULK1 activation, an effect inhibited by activated Akt. ULK1 activation is known to activate ERK1/2 and to be important in cellular survival^{243,297}. Knowing this, we generated mice with a specific microglial deletion of floxed *Ulk1* upon Tamoxifen induced CRE-expression under the CX3CR1 promoter (hereafter referred to as *Ulk1^{fl/fl}Cx3cr1^{CreER}*). In adult mice, the microglia population kinetics was not affected upon this deletion, but after about 2 years, the population was significantly reduced by about 40%. Importantly, this reduction was not compensated by infiltration of other macrophages.


We traced the diminished microglia specifically to a loss of the highly activated ERK1/2 population and that ULK1 deficient microglia suffered from increased ER-stress associated with apoptosis. The loss was most prominent in the cerebral cortex. This chain of autophagy impairment and apoptosis has previously been described in other cells²⁴³. ERK1/2 is suggested to be an upstream regulator of DAM microglia characterized by a reduced CSF1R, increased CSF-1, and further by subpopulation markers such as the MHC class II, TREM2 and CD11C (ITGAX)^{272,298}. In the autophagy-dependent microglia population with highly activated ERK1/2, these markers were found at a higher density, supporting an activated DAM subpopulation attribute.

In order to expand this Autophagy Dependent Age-acquired Microglia, referred to as “ADAM”, we treated aged Wt and *Ulk1^{fl/fl}Cx3cr1^{CreER}* aged mice with intracisternal CSF-1 and IL-34 injections. CSF-1 treatment evoked in both strains a large increase of the microglia less autophagy-dependent and reduced activation of ERK1/2. Conversely, the IL-34 treatment caused a substantial expansion of the ADAM population. This IL-34 induced expansion was specifically canceled in the *Ulk1* deficient mice revealing an aged microglial IL-34/CSF1R/ULK1 axis. Notably, we did not see a significant infiltration of peripheral immune cells in any strain or treatment condition.

The absent ADAM and the reduced microglia population without peripheral infiltration revealed a unique opportunity to study the influence of microglia on disease pathology in the MS model EAE, which is known to aggravate with age. We found the EAE in aged mice deficient in ADAM to associate with high mortality not rescued by IL-34 or CSF-1 treatment even though the latter has similar total microglial counts as aged Wt mice. By quantifying neurons and glial cells by number and apoptosis markers, we found the loss of the ADAM population to associate with a significant reduction of OPC, OLs, and neurons as well as microglia themselves during EAE. Upon IL-34 induced expansion of the ADAM in aged Wt mice, we detected a significant reduction in clinical scores and

increased survival of microglia and CNS cells during EAE. We could also reproduce a neuroprotective effect from IL-34 treated microglia in vitro and detected several neurotrophic factors increased in expression in microglia upon this specific stimulation.

This study discovered and characterized a CNS protective microglial population acquired with age defined by a demand for autophagy protein ULK1 and activated ERK1/2. The absence of this population is associated with aggravated disease phenotypes regardless of total microglial density. Importantly, this population expanded specifically upon IL-34 treatment, a finding to consider when targeting the CSF1R in modeling and treatment of microglia-associated pathologies.



Graphical abstract for Paper II


5.3 Paper III

TGF- β is known to be crucial in establishing a mature microglial homeostatic phenotype^{98,99}. The microglial niche in the CNS parenchyma is, under inflammatory challenges and depletion, populated partly by infiltrating monocytes differentiating into bone marrow-derived macrophages. Our group's previous findings show how depletion is followed by competitive repopulation by microglial cells and monocytes and how these cells adopt shared phenotypes shaped by the tissue surrounding and an intrinsic signature defined by origin⁷⁸. We found the deletion of the TGF- β receptor in *Lyz2* expressing monocytes to associate to reduced integration of bone marrow-derived cells into the CNS parenchyma after microglial depletion. Thus indicating a demand for this cytokine signaling in the migration process or survival and expansion in the target tissue.

In order to silence TGF- β signaling in the repopulated CNS myeloid cells, we utilized the finding of an upregulated *Cx3Cr1* expression in the infiltrating bone-marrow derived cells. This allowed us to employ the inducible deletion of floxed *Tgfb β* ^{fl/fl} upon Tamoxifen induced *Cx3cr1^{CreER}* post infiltration to the CNS of bone-marrow-chimeric mice. The lack

of TGFBR2 in this population altered the transcriptome with upregulation of genes associated with antigen-presentation, inflammation and phagocytosis. About 12 days after gene deletion, the mice developed a progressive motor impairment and died around 10-25 days later. Immunohistochemistry revealed de-myelination and axonal dysfunction in association to myelin-laden giant macrophages in the dorsal spinal cord, causing fore- and hind-limb palsy and late-stage disease also affecting the brain. Without depletion, TGFBR2 deletion specifically in microglia caused similar pathology and symptoms but with later disease-onset compared with the monocyte repopulation model.

In conclusion, we show how bone marrow cells demanded TGF- β signaling to establish a CNS parenchymal myeloid cell and how this population, once settled, requires TGF- β to inhibit a pathogenic macrophage process. This is in line with the need for TGF- β for microglia during developmental establishment and maintenance of a homeostatic phenotype and could be employed in microglial/macrophage replacement therapy and to polarize these populations.



Graphical abstract Paper III

6. DISCUSSION AND FUTURE PERSPECTIVES

This thesis supports the notion of microglia as guardians of the CNS and how malfunctions of these cells cause pathology due to lack of health-promoting capacities. The microglial cell has been defined by localization and morphology for a century, but we still struggle to determine unique microglial signatures and features to distinguish them from bone marrow-derived macrophages. This and the obstacles CNS offer regarding tissue sampling in vivo are factors delaying the understanding of microglia in disease and a health-promoting context.

Microglia-specific gene targeting is still a struggle

The *Cx3Cr1^{CreER}* model targeting microglia is valuable but not completely decisive in its specificity and is likely to be contaminated by BAM. As an alternative, a *Hexb* promotor has been presented as microglia specific and stable in homeostatic and inflammatory conditions²⁹⁹. Another suggestion is to combine a myeloid promoter with a CNS specific, e.g., *Cx3Cr1* and *Sal1* with split Cre-fragments²⁸⁷. However, we should acknowledge the pitfalls of all models in excluding microglial subpopulations and/or including other immune- or CNS-cells.

Microglial depletion and replacement

Microglial replacement therapies upon, e.g., CSF1R inhibition are proposed to ameliorate degenerative and inflammatory neuropathology. Due to the accumulation of leaky mitochondria, impaired autophagy, accumulated mutations, and lipid inclusions, replenished microglia could substantially impact age-associated CNS disease. Our group's previous findings and the results presented in **Paper III** show that bone marrow-derived cells can contribute to a re-established myeloid CNS population following depletion and how TGF- β signaling is demanded and beneficial in maintaining a health-promoting cell in this microglia niche. This can be employed in a desirable treatment regime in polarizing a population that, once established, does not need further pharmaceutical support. **Paper III** further emphasizes previous findings that tell us that microglial origin matters and possibly that replacement therapies should actively inhibit colonization by myeloid cells of bone marrow origin. Given the troubles of depletion models regarding lack of specificity and undesired inflammation caused by cell death and/or toxins, our strategies of specific targeting of repopulating cells presented in **Paper II** and **III** is valuable for future work. In addition to the discovery of myeloid-derived pathogenicity shackled by TGF- β .

Up to date, there is no evidence of bone marrow-derived myeloid contribution to the CNS parenchymal myeloid cell population during homeostasis or even after the acute inflammatory challenge. Yet, we need to study this in a broader context investigating influence from ageing, chronic inflammation, and cytostatic or immune-modulatory treatments since a possible contribution of a bone marrow-derived cell likely affect pathology.

Disease-associated microglia – Loss of a health-promoting phenotype or intrinsic pathogenicity?

Microglia described as “activated”, “amoeboid,” and recently “DAM” have been addressed as pathogenic based on its context of inflammatory and neurodegenerative disease, and

largely by parameters adopted from phenotyping of bone marrow-derived myeloid cells, e.g., molecules associated to antigen-presentation. A likely more relevant idea is now emerging that activated microglia found in the inflammatory CNS lesions have designated functions in restoring a healthy CNS, where phagocytosis is a prominent feature shown in **Paper I** and by others^{4,101,106,134,155,300}. This implies the defects in myelin processing and scavenger receptor recirculation associated with ATG7 deficiency shown in **Paper I** to be assigned a dysfunctional DAM rather than this population to have an intrinsic pathogenic phenotype. It is also possible that the impairment of the key phagocytic function causes an accumulation of this microglial subpopulation, defined by, e.g., CLEC7A density, on the cost of other subpopulations with features important in the recovery from inflammation. However, the DAM phenotypes are, for the most part, defined by single-cell RNA sequencing analyses and still need to be validated as spatially or functionally distinct populations or as transient responsive differentiation stages.

Autophagy-associated phagocytosis in MS and EAE

The list of suggested diseases with myeloid cell autophagy-associated phagocytic dysfunction is growing, and this pathway cannot be viewed as something irregular. One question that remains unanswered is if this is only a degradation pathway, or if the cargo selection has specific regulatory functions, and if the cargo by vesicle specifics is destined towards certain functions, e.g., antigen presentation. Another issue is how activation of this RUBICON-directed pathway affects canonical-autophagy functions during long-term exposure to, e.g., myelin or apoptotic cells. The process of autophagy-associated phagocytosis is also largely understudied in human disease, and findings by us in **Paper I** and others need to be reproduced or supported by studies on the pathogenesis of the human disease. This is also called for by the autophagy-associated genes and pathways detected in MS susceptibility genetics.

CSF-1 vs. IL-34 promoted microglial subpopulations in the aged CNS

The aged CNS was further explored in **Paper II** where we found a microglial population with activated ERK1/2, a known upstream regulator of the DAM microglia phenotype, to be dependent on canonical autophagy. This population referred to as “ADAM” was found protective during EAE, and loss of this population in the aged mice deficient in microglial ULK1 associated with high mortality, immune cell infiltration, and CNS cell death. Recent publications have explored the regional developmental disparate effects of IL-34 and CSF-1. We could expand the ADAM population specifically upon IL-34 treatment, which significantly reduced the signs of neuroinflammation compared to the CSF-1 treatment, which selectively expanded the autophagy-independent population with low ERK1/2 activation. We should also highlight that even though CSF-1 is associated with DAM and act on other cells than microglia, this treatment did not aggravate EAE disease. The evidently increased density of activation markers in the IL-34 expanded ADAM population supports a microglial DAM-phenotype with protective capacity. However, besides the IL-34 induced expression of neurotrophic factors the executive functions derived specifically from the two CSF1R ligands and to which degree they are redundant remains to be shown.

In conclusion, **Paper II** highlights the demand for canonical autophagy to maintain a microglial population with unique neuroprotective features during EAE. We can also view this model as a partial microglial depletion model, where the depletion in comparison to

other models is less effective but silent and not compensated by bone marrow-derived myeloid cells. How this affects the ageing CNS could be explored further in homeostatic conditions or other disease models, including exploring the suggested IL-34 specific effects.

The findings in **Paper II** revealing the age-acquired microglial subpopulation with higher demand for canonical-autophagy, suggesting this pathway rather more activated than declined with age. This is somewhat supported by an altered microglial phenotype downstream mTOR signaling recently reported²³⁴. However, in **Paper I**, we find an increased expression of DAM-associated genes in microglia deficient in both canonical and non-canonical autophagy shared partly with aged microglia. This further supports that the autophagy-dependent phenotype or subpopulation is acquired during ageing. One could also speculate an increased canonical-autophagy to occupy the autophagy proteins in aged microglia and thereby reduce the capacity for non-canonical autophagy associated phagocytosis — a deficiency we targeted by trehalose treatment in **Paper I**.

Induced autophagy as a potential therapeutic target in neuroinflammation

Loss of EAE recovery in the C57BL/6 model should probably be addressed as a prolonged inflammatory relapse rather than a transition to the progressive phenotype seen in human MS. This does not exclude shared specific pathways such as microglial autophagy. It is also possible that the aggravated EAE associated with age suffers from the same pathology as the progressive human disease. In contrast to models of neurodegeneration, studies in EAE addressing age is underutilized and likely informative.

The phenotypes presented in **Paper I** and **II** add to an emerging concept of autophagy as regulatory and executive function in myeloid cells. While immune-modulatory treatments commonly dampen the immune surveillance at large, autophagy induction tentatively has beneficial effects in surveillance of tumor cells and microbes and simultaneously dampening an aberrant immune activation. In general, boosting an immune function is rare in comparison to restraining immunity as a treatment regime. If this is due to complications or is an underused path remains to be seen.

The findings of trehalose ameliorating age-associated neuroinflammation in **Paper I** are promising and novel in targeting autophagy-associated phagocytosis. This could potentially be implemented as a dietary adjustment to reduce the risk of developing a disease or halt disease progression. However, deliberate trehalose supplementation should also be considered an option.

In this thesis, I present findings and ideas of health-promoting pathways in cells of the CNS microglial niche showing cellular phenotypes and functional outcomes. Knowledge of these cells is rapidly expanding and holds answers and offers solutions redeeming neuroinflammation that has been blindsided by methodological challenges.

7. ACKNOWLEDGMENTS

What a slow, busy, and dazzling journey this has been. Like an Indian train ride. I'm thankful for the possibility to do research at the kind and encouraging **Neuroimmunology unit** at **Karolinska Institutet/Centrum for molecular medicine**. I also want to thank the **Karolinska Institutet** for funding and first-class education the past 15 years.

To my wise and visionary main supervisor **Tomas**, thank you for your sharp perspectives and for always being positive and encouraging. And for bringing in the \$\$\$\$. To my cool and intelligent co-supervisor, **Maja**, thank you for asking the relevant questions at the right time, and your quality standards. And to **Andre**, my charismatic and clever co-supervisor and friend. Thank you for taking the time (almost every day) to untangle many practical and intellectual matters.

I'm so grateful for all that I have learned you all, and for the efforts in sharpening the narratives.

I arrived in the **Neuroimmunology unit** close to a decade ago. It was autumn, and I was a bit confused. The welcome I got from the "**old rat-gang**" was very warm and inclusive. From that era, I would like to address a few (of many) inspiring friends:

Melanie – Thank you for your sunny smartness and a perfect introduction to scientific labor and for handing over the "autophagy project", which still is my *Sagrada familia*.

Roham, thank you for your inspiring next-level aspirations and competence. You gave me valuable advice already from the start ("If you don't work during easter, you're not a real scientist", and then you....). **Petra** – I remember at first you tried to be scary so that I wouldn't mess up the "RNAse free room". Nevertheless, I think I did, and you are still my friend, sweetheart, and such a skilled scientist.

And to the rest of the gang: **Ame, Shahin, Hannes, Nada, Alan, Maria, Pernilla, Sevi** - Thank you for all the fun times and discussions!

Then came a couple of new kids to the block (I'm so glad to now call you and your families my close friends). **Karl** – It took 8 minutes to sense a good friendship coming. Because of your somewhat confusing ironic description of things, it took a little bit longer to see what a great scientist you are. **Harald** – I would be lost in the maze of myeloid cells without you. Many happy memories, not least during the traveling escapades, spreading the word of microglia.

Thank you to **Bob** for taking an interest in me, for the nice talks, and to your talented and helpful team of "big-eater" experts - **Melanie, Keying, Sohel, Jinming, Xingmei** (please never stop laughing). To **Fredrik, Ingrid, and Lou** for expertise and support, and to your excellent team-members **Faiez, Cecilia, Richard, Micke, Magda, Tojo, Rux, Mathias, and Susi**. And to **Marie** – with the best of forwardness and spark. To **Sabrina** – your devotion, helpfulness, and technical skills are gold. To **Ewoud, Majid, Eliane, and Lara** for being bright and fun, and for your scientific assistance and discussions. To **Manuel** and **Milena** for great and educative collaborative work. And to **Nånnis** for friendship, inspiration, and sharing views on clinical/research-work dynamics.

I am grateful to **Helena, Jan-Alvar, the other organizers**, and the participants of the research school **NCRSCID**. Thanks for the excellent education and widening of my research vision. And to professor **Katja Simon**, Kennedy institute Oxford, who guided me through the autophagy pathways at critical moments during my Ph.D.

And **Mohsen**. You are the caretaking enigma of this fine establishment. Thanks for everything! A special thanks also to the animal facility personnel, especially **Helene Kungsmark**, for being so deer and helpful. To **Brinda, Venus**, and **Gun** – thanks for keeping the laboratory wheels spinning. And my warmest gratitude to all the collaborators, funding bodies, current and previous neuroimmunology and rheumatology unit members this past decade not mentioned by name.

Thanks to co-workers at the **Karolinska Huddinge hospital emergency care units**, your extra efforts fuel the health-care system, and the tough and emotional work during the pandemic should be honored. At least with a song and a statue.

To my inspiring **mom, pops, sisters, brother, uncles, aunts, grandparents**, and **cousins** - The spirit you have is the soil of this work. To old and new **friends**. And to **Barnens Ö** for providing a beautiful escape during the summers these past years.

Last but foremost, I would like to thank my beloved family - **Moa, Edgar, Elis**, and **Emmylou** – Thanks for all the support in this challenging and world of science, which sometimes is hard to explain. And the needed happy diversions.

I'm glad I've had the chance to thrive with the support of this community to transform my ideas into something to be proud of. I hope this is not the last I see of you **KI Neuroimmunology people**.

Until then.

Peace /Rasmus

8. REFERENCES

- 1 Walton, C. *et al.* Rising prevalence of multiple sclerosis worldwide: Insights from the Atlas of MS, third edition. *Multiple sclerosis (Houndmills, Basingstoke, England)*, 1352458520970841, doi:10.1177/1352458520970841 (2020).
- 2 Global, regional, and national burden of neurological disorders, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet. Neurology* **18**, 459-480, doi:10.1016/s1474-4422(18)30499-x (2019).
- 3 Filippi, M. *et al.* Multiple sclerosis. *Nature Reviews Disease Primers* **4**, 43, doi:10.1038/s41572-018-0041-4 (2018).
- 4 Deczkowska, A. *et al.* Disease-Associated Microglia: A Universal Immune Sensor of Neurodegeneration. *Cell* **173**, 1073-1081, doi:10.1016/j.cell.2018.05.003 (2018).
- 5 Friese, M. A., Schattling, B. & Fugger, L. Mechanisms of neurodegeneration and axonal dysfunction in multiple sclerosis. *Nat. Rev. Neurol.* **10**, 225-238, doi:10.1038/nrneurol.2014.37 (2014).
- 6 Schattling, B. *et al.* Bassoon proteinopathy drives neurodegeneration in multiple sclerosis. *Nature neuroscience* **22**, 887-896, doi:10.1038/s41593-019-0385-4 (2019).
- 7 Multiple sclerosis genomic map implicates peripheral immune cells and microglia in susceptibility. *Science (New York, N.Y.)* **365**, doi:10.1126/science.aav7188 (2019).
- 8 Olsson, T., Barcellos, L. F. & Alfredsson, L. Interactions between genetic, lifestyle and environmental risk factors for multiple sclerosis. *Nature reviews. Neurology* **13**, 25-36, doi:10.1038/nrneurol.2016.187 (2017).
- 9 van Langelaar, J., Rijvers, L., Smolders, J. & van Luijn, M. M. B and T Cells Driving Multiple Sclerosis: Identity, Mechanisms and Potential Triggers. *Frontiers in immunology* **11**, doi:10.3389/fimmu.2020.00760 (2020).
- 10 Jelcic, I. *et al.* Memory B Cells Activate Brain-Homing, Autoreactive CD4(+) T Cells in Multiple Sclerosis. *Cell* **175**, 85-100.e123, doi:10.1016/j.cell.2018.08.011 (2018).
- 11 Hauser, S. L. *et al.* Ocrelizumab versus Interferon Beta-1a in Relapsing Multiple Sclerosis. *The New England journal of medicine* **376**, 221-234, doi:10.1056/NEJMoa1601277 (2017).
- 12 Polman, C. H. *et al.* A randomized, placebo-controlled trial of natalizumab for relapsing multiple sclerosis. *The New England journal of medicine* **354**, 899-910, doi:10.1056/NEJMoa044397 (2006).
- 13 Zrzavy, T. *et al.* Loss of 'homeostatic' microglia and patterns of their activation in active multiple sclerosis. *Brain : a journal of neurology* **140**, 1900-1913, doi:10.1093/brain/awx113 (2017).
- 14 Bai, Z. *et al.* Cerebrospinal Fluid and Blood Cytokines as Biomarkers for Multiple Sclerosis: A Systematic Review and Meta-Analysis of 226 Studies With 13,526 Multiple Sclerosis Patients. *Front Neurosci* **13**, 1026-1026, doi:10.3389/fnins.2019.01026 (2019).
- 15 Restorick, S. M. *et al.* CCR6(+) Th cells in the cerebrospinal fluid of persons with multiple sclerosis are dominated by pathogenic non-classic Th1 cells and GM-CSF-only-secreting Th cells. *Brain, behavior, and immunity* **64**, 71-79, doi:10.1016/j.bbi.2017.03.008 (2017).
- 16 Khademi, M. *et al.* Cerebrospinal fluid CXCL13 in multiple sclerosis: a suggestive prognostic marker for the disease course. *Multiple sclerosis (Houndmills, Basingstoke, England)* **17**, 335-343, doi:10.1177/1352458510389102 (2011).

- 17 Bagaeva, L. V., Rao, P., Powers, J. M. & Segal, B. M. CXC Chemokine Ligand 13 Plays a Role in Experimental Autoimmune Encephalomyelitis. *The Journal of Immunology* **176**, 7676-7685, doi:10.4049/jimmunol.176.12.7676 (2006).
- 18 Croxford, A. L. *et al.* The Cytokine GM-CSF Drives the Inflammatory Signature of CCR2⁺ Monocytes and Licenses Autoimmunity. *Immunity* **43**, 502-514, doi:10.1016/j.immuni.2015.08.010 (2015).
- 19 Ronchi, F. *et al.* Experimental priming of encephalitogenic Th1/Th17 cells requires pertussis toxin-driven IL-1 β production by myeloid cells. *Nature communications* **7**, 11541, doi:10.1038/ncomms11541 (2016).
- 20 Spath, S. *et al.* Dysregulation of the Cytokine GM-CSF Induces Spontaneous Phagocyte Invasion and Immunopathology in the Central Nervous System. *Immunity* **46**, 245-260, doi:10.1016/j.immuni.2017.01.007 (2017).
- 21 Vogel, D. Y. *et al.* GM-CSF promotes migration of human monocytes across the blood brain barrier. *European journal of immunology* **45**, 1808-1819, doi:10.1002/eji.201444960 (2015).
- 22 McQualter, J. L. *et al.* Granulocyte macrophage colony-stimulating factor: a new putative therapeutic target in multiple sclerosis. *The Journal of experimental medicine* **194**, 873-882, doi:10.1084/jem.194.7.873 (2001).
- 23 Paré, A. *et al.* IL-1 β enables CNS access to CCR2^{hi} monocytes and the generation of pathogenic cells through GM-CSF released by CNS endothelial cells. *Proceedings of the National Academy of Sciences* **115**, E1194-E1203, doi:10.1073/pnas.1714948115 (2018).
- 24 Becher, B., Durell, B. G. & Noelle, R. J. IL-23 produced by CNS-resident cells controls T cell encephalitogenicity during the effector phase of experimental autoimmune encephalomyelitis. *The Journal of clinical investigation* **112**, 1186-1191, doi:10.1172/jci19079 (2003).
- 25 Arellano, G., Ottum, P. A., Reyes, L. I., Burgos, P. I. & Naves, R. Stage-Specific Role of Interferon-Gamma in Experimental Autoimmune Encephalomyelitis and Multiple Sclerosis. *Frontiers in immunology* **6**, 492, doi:10.3389/fimmu.2015.00492 (2015).
- 26 Cua, D. J. *et al.* Interleukin-23 rather than interleukin-12 is the critical cytokine for autoimmune inflammation of the brain. *Nature* **421**, 744-748, doi:10.1038/nature01355 (2003).
- 27 Komiyama, Y. *et al.* IL-17 plays an important role in the development of experimental autoimmune encephalomyelitis. *Journal of immunology (Baltimore, Md. : 1950)* **177**, 566-573, doi:10.4049/jimmunol.177.1.566 (2006).
- 28 Mahad, D. H., Trapp, B. D. & Lassmann, H. Pathological mechanisms in progressive multiple sclerosis. *The Lancet. Neurology* **14**, 183-193, doi:10.1016/s1474-4422(14)70256-x (2015).
- 29 Lassmann, H. Targets of therapy in progressive MS. *Multiple sclerosis (Houndmills, Basingstoke, England)* **23**, 1593-1599, doi:10.1177/1352458517729455 (2017).
- 30 Lassmann, H. Cortical lesions in multiple sclerosis: inflammation versus neurodegeneration. *Brain : a journal of neurology* **135**, 2904-2905, doi:10.1093/brain/aws260 (2012).
- 31 Machado-Santos, J. *et al.* The compartmentalized inflammatory response in the multiple sclerosis brain is composed of tissue-resident CD8⁺ T lymphocytes and B cells. *Brain : a journal of neurology* **141**, 2066-2082, doi:10.1093/brain/awy151 (2018).

- 32 Luchetti, S. *et al.* Progressive multiple sclerosis patients show substantial lesion activity that correlates with clinical disease severity and sex: a retrospective autopsy cohort analysis. *Acta Neuropathologica* **135**, 511-528, doi:10.1007/s00401-018-1818-y (2018).
- 33 O'Loughlin, E., Madore, C., Lassmann, H. & Butovsky, O. Microglial Phenotypes and Functions in Multiple Sclerosis. *Cold Spring Harbor perspectives in medicine* **8**, a028993, doi:10.1101/cshperspect.a028993 (2018).
- 34 van der Poel, M. *et al.* Transcriptional profiling of human microglia reveals grey-white matter heterogeneity and multiple sclerosis-associated changes. *Nature communications* **10**, 1139, doi:10.1038/s41467-019-08976-7 (2019).
- 35 Correale, J., Gaitán, M. I., Ysrraelit, M. C. & Fiol, M. P. Progressive multiple sclerosis: from pathogenic mechanisms to treatment. *Brain : a journal of neurology* **140**, 527-546, doi:10.1093/brain/aww258 (2017).
- 36 Constantinescu, C. S., Farooqi, N., O'Brien, K. & Gran, B. Experimental autoimmune encephalomyelitis (EAE) as a model for multiple sclerosis (MS). *British journal of pharmacology* **164**, 1079-1106, doi:10.1111/j.1476-5381.2011.01302.x (2011).
- 37 Rivers, T. M., Sprunt, D. H. & Berry, G. P. OBSERVATIONS ON ATTEMPTS TO PRODUCE ACUTE DISSEMINATED ENCEPHALOMYELITIS IN MONKEYS. *The Journal of experimental medicine* **58**, 39-53, doi:10.1084/jem.58.1.39 (1933).
- 38 Lassmann, H. & Bradl, M. Multiple sclerosis: experimental models and reality. *Acta neuropathologica* **133**, 223-244, doi:10.1007/s00401-016-1631-4 (2017).
- 39 Friese, M. A. *et al.* Opposing effects of HLA class I molecules in tuning autoreactive CD8+ T cells in multiple sclerosis. *Nature medicine* **14**, 1227-1235, doi:10.1038/nm.1881 (2008).
- 40 Sun, D. *et al.* Myelin antigen-specific CD8+ T cells are encephalitogenic and produce severe disease in C57BL/6 mice. *Journal of immunology (Baltimore, Md. : 1950)* **166**, 7579-7587, doi:10.4049/jimmunol.166.12.7579 (2001).
- 41 Wagner, C. A., Roqué, P. J., Mileur, T. R., Liggitt, D. & Goverman, J. M. Myelin-specific CD8+ T cells exacerbate brain inflammation in CNS autoimmunity. *The Journal of clinical investigation* **130**, 203-213, doi:10.1172/jci132531 (2020).
- 42 Saxena, A. *et al.* Cutting edge: Multiple sclerosis-like lesions induced by effector CD8 T cells recognizing a sequestered antigen on oligodendrocytes. *Journal of immunology (Baltimore, Md. : 1950)* **181**, 1617-1621, doi:10.4049/jimmunol.181.3.1617 (2008).
- 43 Weber, M. S. *et al.* B-cell activation influences T-cell polarization and outcome of anti-CD20 B-cell depletion in central nervous system autoimmunity. *Annals of neurology* **68**, 369-383, doi:10.1002/ana.22081 (2010).
- 44 Matsushita, T., Yanaba, K., Bouaziz, J. D., Fujimoto, M. & Tedder, T. F. Regulatory B cells inhibit EAE initiation in mice while other B cells promote disease progression. *The Journal of clinical investigation* **118**, 3420-3430, doi:10.1172/jci36030 (2008).
- 45 McCarthy, D. P., Richards, M. H. & Miller, S. D. Mouse models of multiple sclerosis: experimental autoimmune encephalomyelitis and Theiler's virus-induced demyelinating disease. *Methods in molecular biology (Clifton, N.J.)* **900**, 381-401, doi:10.1007/978-1-60761-720-4_19 (2012).
- 46 Osorio-Querejeta, I., Sáenz-Cuesta, M., Muñoz-Culla, M. & Otaegui, D. Models for Studying Myelination, Demyelination and Remyelination. *NeuroMolecular Medicine* **19**, 181-192, doi:10.1007/s12017-017-8442-1 (2017).

- 47 Ginhoux, F., Lim, S., Hoeffel, G., Low, D. & Huber, T. Origin and differentiation of microglia. *Frontiers in cellular neuroscience* **7**, 45, doi:10.3389/fncel.2013.00045 (2013).
- 48 Schulz, C. *et al.* A lineage of myeloid cells independent of Myb and hematopoietic stem cells. *Science (New York, N.Y.)* **336**, 86-90, doi:10.1126/science.1219179 (2012).
- 49 Ginhoux, F. *et al.* Fate mapping analysis reveals that adult microglia derive from primitive macrophages. *Science (New York, N.Y.)* **330**, 841-845, doi:10.1126/science.1194637 (2010).
- 50 Chen, H. R. *et al.* Fate mapping via CCR2-CreER mice reveals monocyte-to-microglia transition in development and neonatal stroke. *Science advances* **6**, eabb2119, doi:10.1126/sciadv.abb2119 (2020).
- 51 Réu, P. *et al.* The Lifespan and Turnover of Microglia in the Human Brain. *Cell reports* **20**, 779-784, doi:10.1016/j.celrep.2017.07.004 (2017).
- 52 Askew, K. *et al.* Coupled Proliferation and Apoptosis Maintain the Rapid Turnover of Microglia in the Adult Brain. *Cell reports* **18**, 391-405, doi:10.1016/j.celrep.2016.12.041 (2017).
- 53 Kierdorf, K. *et al.* Microglia emerge from erythromyeloid precursors via Pu.1- and Irf8-dependent pathways. *Nature neuroscience* **16**, 273-280, doi:10.1038/nn.3318 (2013).
- 54 Erblich, B., Zhu, L., Etgen, A. M., Dobrenis, K. & Pollard, J. W. Absence of colony stimulation factor-1 receptor results in loss of microglia, disrupted brain development and olfactory deficits. *PloS one* **6**, e26317, doi:10.1371/journal.pone.0026317 (2011).
- 55 Oosterhof, N. *et al.* Homozygous Mutations in CSF1R Cause a Pediatric-Onset Leukoencephalopathy and Can Result in Congenital Absence of Microglia. *American journal of human genetics* **104**, 936-947, doi:10.1016/j.ajhg.2019.03.010 (2019).
- 56 Easley-Neal, C., Foreman, O., Sharma, N., Zarrin, A. A. & Weimer, R. M. CSF1R Ligands IL-34 and CSF1 Are Differentially Required for Microglia Development and Maintenance in White and Gray Matter Brain Regions. *Frontiers in immunology* **10**, 2199, doi:10.3389/fimmu.2019.02199 (2019).
- 57 Obst, J. *et al.* Inhibition of IL-34 Unveils Tissue-Selectivity and Is Sufficient to Reduce Microglial Proliferation in a Model of Chronic Neurodegeneration. *Frontiers in immunology* **11**, doi:10.3389/fimmu.2020.579000 (2020).
- 58 Wei, S. *et al.* Functional overlap but differential expression of CSF-1 and IL-34 in their CSF-1 receptor-mediated regulation of myeloid cells. *Journal of leukocyte biology* **88**, 495-505, doi:10.1189/jlb.1209822 (2010).
- 59 Baghdadi, M. *et al.* Interleukin-34, a comprehensive review. *Journal of leukocyte biology* **104**, 931-951, doi:10.1002/jlb.Mr1117-457r (2018).
- 60 Zhang, Y. *et al.* An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **34**, 11929-11947, doi:10.1523/jneurosci.1860-14.2014 (2014).
- 61 Zhang, Y. *et al.* Purification and Characterization of Progenitor and Mature Human Astrocytes Reveals Transcriptional and Functional Differences with Mouse. *Neuron* **89**, 37-53, doi:10.1016/j.neuron.2015.11.013 (2016).

- 62 Fernández-Calle, R. *et al.* Role of RPTP β/ζ in neuroinflammation and microglia-neuron communication. *Scientific reports* **10**, 20259, doi:10.1038/s41598-020-76415-5 (2020).
- 63 Kana, V. *et al.* CSF-1 controls cerebellar microglia and is required for motor function and social interaction. *The Journal of experimental medicine* **216**, 2265-2281, doi:10.1084/jem.20182037 (2019).
- 64 Greter, M. *et al.* Stroma-derived interleukin-34 controls the development and maintenance of langerhans cells and the maintenance of microglia. *Immunity* **37**, 1050-1060, doi:10.1016/j.immuni.2012.11.001 (2012).
- 65 Kondo, Y. & Duncan, I. D. Selective reduction in microglia density and function in the white matter of colony-stimulating factor-1-deficient mice. *Journal of neuroscience research* **87**, 2686-2695, doi:10.1002/jnr.22096 (2009).
- 66 Stanley, E. R. & Chitu, V. CSF-1 receptor signaling in myeloid cells. *Cold Spring Harbor perspectives in biology* **6**, doi:10.1101/cshperspect.a021857 (2014).
- 67 Richardson, E. T. *et al.* ERK Signaling Is Essential for Macrophage Development. *PloS one* **10**, e0140064, doi:10.1371/journal.pone.0140064 (2015).
- 68 Tay, T. L. *et al.* A new fate mapping system reveals context-dependent random or clonal expansion of microglia. *Nature neuroscience* **20**, 793-803, doi:10.1038/nn.4547 (2017).
- 69 Ajami, B., Bennett, J. L., Krieger, C., Tetzlaff, W. & Rossi, F. M. Local self-renewal can sustain CNS microglia maintenance and function throughout adult life. *Nature neuroscience* **10**, 1538-1543, doi:10.1038/nn2014 (2007).
- 70 Huang, Y. *et al.* Repopulated microglia are solely derived from the proliferation of residual microglia after acute depletion. *Nature neuroscience* **21**, 530-540, doi:10.1038/s41593-018-0090-8 (2018).
- 71 Tan, Y.-L., Yuan, Y. & Tian, L. Microglial regional heterogeneity and its role in the brain. *Molecular Psychiatry* **25**, 351-367, doi:10.1038/s41380-019-0609-8 (2020).
- 72 Masuda, T. *et al.* Spatial and temporal heterogeneity of mouse and human microglia at single-cell resolution. *Nature* **566**, 388-392, doi:10.1038/s41586-019-0924-x (2019).
- 73 Lawson, L. J., Perry, V. H., Dri, P. & Gordon, S. Heterogeneity in the distribution and morphology of microglia in the normal adult mouse brain. *Neuroscience* **39**, 151-170, doi:10.1016/0306-4522(90)90229-w (1990).
- 74 Easley-Neal, C., Foreman, O., Sharma, N., Zarrin, A. A. & Weimer, R. M. CSF1R Ligands IL-34 and CSF1 Are Differentially Required for Microglia Development and Maintenance in White and Gray Matter Brain Regions. *Frontiers in immunology* **10**, doi:10.3389/fimmu.2019.02199 (2019).
- 75 Boulakirba, S. *et al.* IL-34 and CSF-1 display an equivalent macrophage differentiation ability but a different polarization potential. *Scientific reports* **8**, 256, doi:10.1038/s41598-017-18433-4 (2018).
- 76 Wlodarczyk, A. *et al.* CSF1R Stimulation Promotes Increased Neuroprotection by CD11c+ Microglia in EAE. *Frontiers in cellular neuroscience* **12**, doi:10.3389/fncel.2018.00523 (2019).
- 77 O'Neil, S. M., Witcher, K. G., McKim, D. B. & Godbout, J. P. Forced turnover of aged microglia induces an intermediate phenotype but does not rebalance CNS environmental cues driving priming to immune challenge. *Acta neuropathologica communications* **6**, 129, doi:10.1186/s40478-018-0636-8 (2018).

- 78 Lund, H. *et al.* Competitive repopulation of an empty microglial niche yields functionally distinct subsets of microglia-like cells. *Nature communications* **9**, 4845, doi:10.1038/s41467-018-07295-7 (2018).
- 79 Cronk, J. C. *et al.* Peripherally derived macrophages can engraft the brain independent of irradiation and maintain an identity distinct from microglia. *The Journal of experimental medicine* **215**, 1627-1647, doi:10.1084/jem.20180247 (2018).
- 80 Varvel, N. H. *et al.* Microglial repopulation model reveals a robust homeostatic process for replacing CNS myeloid cells. *Proceedings of the National Academy of Sciences of the United States of America* **109**, 18150-18155, doi:10.1073/pnas.1210150109 (2012).
- 81 Han, J., Harris, R. A. & Zhang, X.-M. An updated assessment of microglia depletion: current concepts and future directions. *Molecular Brain* **10**, 25, doi:10.1186/s13041-017-0307-x (2017).
- 82 Huang, Y. *et al.* Repopulated microglia are solely derived from the proliferation of residual microglia after acute depletion. *Nature neuroscience* **21**, 530-540, doi:10.1038/s41593-018-0090-8 (2018).
- 83 Agudo, J. *et al.* GFP-specific CD8 T cells enable targeted cell depletion and visualization of T-cell interactions. *Nature biotechnology* **33**, 1287-1292, doi:10.1038/nbt.3386 (2015).
- 84 Bruttger, J. *et al.* Genetic Cell Ablation Reveals Clusters of Local Self-Renewing Microglia in the Mammalian Central Nervous System. *Immunity* **43**, 92-106, doi:10.1016/j.immuni.2015.06.012 (2015).
- 85 Lei, F. *et al.* CSF1R inhibition by a small-molecule inhibitor is not microglia specific; affecting hematopoiesis and the function of macrophages. *Proceedings of the National Academy of Sciences* **117**, 23336-23338, doi:10.1073/pnas.1922788117 (2020).
- 86 Tanabe, S., Saitoh, S., Miyajima, H., Itokazu, T. & Yamashita, T. Microglia suppress the secondary progression of autoimmune encephalomyelitis. *Glia* **67**, 1694-1704, doi:10.1002/glia.23640 (2019).
- 87 Sariol, A. *et al.* Microglia depletion exacerbates demyelination and impairs remyelination in a neurotropic coronavirus infection. *Proceedings of the National Academy of Sciences* **117**, 24464-24474, doi:10.1073/pnas.2007814117 (2020).
- 88 Kotter, M. R., Zhao, C., van Rooijen, N. & Franklin, R. J. Macrophage-depletion induced impairment of experimental CNS remyelination is associated with a reduced oligodendrocyte progenitor cell response and altered growth factor expression. *Neurobiology of disease* **18**, 166-175, doi:10.1016/j.nbd.2004.09.019 (2005).
- 89 Zhu, K. *et al.* Absence of microglia or presence of peripherally-derived macrophages does not affect tau pathology in young or old hTau mice. *Glia* **68**, 1466-1478, doi:https://doi.org/10.1002/glia.23794 (2020).
- 90 Beckmann, N. *et al.* Brain region-specific enhancement of remyelination and prevention of demyelination by the CSF1R kinase inhibitor BLZ945. *Acta neuropathologica communications* **6**, 9, doi:10.1186/s40478-018-0510-8 (2018).
- 91 Spangenberg, E. *et al.* Sustained microglial depletion with CSF1R inhibitor impairs parenchymal plaque development in an Alzheimer's disease model. *Nature communications* **10**, 3758, doi:10.1038/s41467-019-11674-z (2019).

- 92 Olmos-Alonso, A. *et al.* Pharmacological targeting of CSF1R inhibits microglial proliferation and prevents the progression of Alzheimer's-like pathology. *Brain : a journal of neurology* **139**, 891-907, doi:10.1093/brain/awv379 (2016).
- 93 Kierdorf, K. & Prinz, M. Microglia in steady state. *The Journal of clinical investigation* **127**, 3201-3209, doi:10.1172/jci90602 (2017).
- 94 Krasemann, S. *et al.* The TREM2-APOE Pathway Drives the Transcriptional Phenotype of Dysfunctional Microglia in Neurodegenerative Diseases. *Immunity* **47**, 566-581.e569, doi:10.1016/j.immuni.2017.08.008 (2017).
- 95 Böttcher, C. *et al.* Human microglia regional heterogeneity and phenotypes determined by multiplexed single-cell mass cytometry. *Nature neuroscience* **22**, 78-90, doi:10.1038/s41593-018-0290-2 (2019).
- 96 Sankowski, R. *et al.* Mapping microglia states in the human brain through the integration of high-dimensional techniques. *Nature neuroscience* **22**, 2098-2110, doi:10.1038/s41593-019-0532-y (2019).
- 97 Hammond, T. R. *et al.* Single-Cell RNA Sequencing of Microglia throughout the Mouse Lifespan and in the Injured Brain Reveals Complex Cell-State Changes. *Immunity* **50**, 253-271.e256, doi:10.1016/j.immuni.2018.11.004 (2019).
- 98 Butovsky, O. *et al.* Identification of a unique TGF- β -dependent molecular and functional signature in microglia. *Nature neuroscience* **17**, 131-143, doi:10.1038/nn.3599 (2014).
- 99 Zöller, T. *et al.* Silencing of TGF β signalling in microglia results in impaired homeostasis. *Nature communications* **9**, 4011, doi:10.1038/s41467-018-06224-y (2018).
- 100 Butovsky, O. *et al.* Identification of a unique TGF- β -dependent molecular and functional signature in microglia. *Nature neuroscience* **17**, 131-143, doi:10.1038/nn.3599 (2014).
- 101 Olah, M. *et al.* Identification of a microglia phenotype supportive of remyelination. *Glia* **60**, 306-321, doi:10.1002/glia.21266 (2012).
- 102 Lalive, P. H. *et al.* TGF-beta-treated microglia induce oligodendrocyte precursor cell chemotaxis through the HGF-c-Met pathway. *European journal of immunology* **35**, 727-737, doi:10.1002/eji.200425430 (2005).
- 103 Abutbul, S. *et al.* TGF- β signaling through SMAD2/3 induces the quiescent microglial phenotype within the CNS environment. *Glia* **60**, 1160-1171, doi:10.1002/glia.22343 (2012).
- 104 Goldmann, T. *et al.* A new type of microglia gene targeting shows TAK1 to be pivotal in CNS autoimmune inflammation. *Nature neuroscience* **16**, 1618-1626, doi:10.1038/nn.3531 (2013).
- 105 Keren-Shaul, H. *et al.* A Unique Microglia Type Associated with Restricting Development of Alzheimer's Disease. *Cell* **169**, 1276-1290.e1217, doi:10.1016/j.cell.2017.05.018 (2017).
- 106 Jordão, M. J. C. *et al.* Single-cell profiling identifies myeloid cell subsets with distinct fates during neuroinflammation. *Science (New York, N.Y.)* **363**, doi:10.1126/science.aat7554 (2019).
- 107 Lloyd, A. F. & Miron, V. E. The pro-remyelination properties of microglia in the central nervous system. *Nat. Rev. Neurol.* **15**, 447-458, doi:10.1038/s41582-019-0184-2 (2019).
- 108 Park, C. *et al.* The landscape of myeloid and astrocyte phenotypes in acute multiple sclerosis lesions. *Acta neuropathologica communications* **7**, 130, doi:10.1186/s40478-019-0779-2 (2019).

- 109 Fischer, M. T. *et al.* NADPH oxidase expression in active multiple sclerosis lesions in relation to oxidative tissue damage and mitochondrial injury. *Brain : a journal of neurology* **135**, 886-899, doi:10.1093/brain/aws012 (2012).
- 110 Binder, M. D. *et al.* Gas6 deficiency increases oligodendrocyte loss and microglial activation in response to cuprizone-induced demyelination. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **28**, 5195-5206, doi:10.1523/jneurosci.1180-08.2008 (2008).
- 111 Cantoni, C. *et al.* TREM2 regulates microglial cell activation in response to demyelination in vivo. *Acta Neuropathol* **129**, 429-447, doi:10.1007/s00401-015-1388-1 (2015).
- 112 Lampron, A. *et al.* Inefficient clearance of myelin debris by microglia impairs remyelinating processes. *The Journal of experimental medicine* **212**, 481-495, doi:10.1084/jem.20141656 (2015).
- 113 Kotter, M. R., Li, W. W., Zhao, C. & Franklin, R. J. Myelin impairs CNS remyelination by inhibiting oligodendrocyte precursor cell differentiation. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **26**, 328-332, doi:10.1523/jneurosci.2615-05.2006 (2006).
- 114 Yamasaki, R. *et al.* Differential roles of microglia and monocytes in the inflamed central nervous system. *The Journal of experimental medicine* **211**, 1533-1549, doi:10.1084/jem.20132477 (2014).
- 115 Dendrou, C. A., Fugger, L. & Friese, M. A. Immunopathology of multiple sclerosis. *Nat. Rev. Immunol.* **15**, 545-558, doi:10.1038/nri3871 (2015).
- 116 Bramow, S. *et al.* Demyelination versus remyelination in progressive multiple sclerosis. *Brain : a journal of neurology* **133**, 2983-2998, doi:10.1093/brain/awq250 (2010).
- 117 Lubetzki, C., Zalc, B., Williams, A., Stadelmann, C. & Stankoff, B. Remyelination in multiple sclerosis: from basic science to clinical translation. *The Lancet. Neurology* **19**, 678-688, doi:10.1016/s1474-4422(20)30140-x (2020).
- 118 Guo, L. *et al.* Bi-allelic CSF1R Mutations Cause Skeletal Dysplasia of Dysosteosclerosis-Pyle Disease Spectrum and Degenerative Encephalopathy with Brain Malformation. *American journal of human genetics* **104**, 925-935, doi:10.1016/j.ajhg.2019.03.004 (2019).
- 119 Miron, V. E. *et al.* M2 microglia and macrophages drive oligodendrocyte differentiation during CNS remyelination. *Nature neuroscience* **16**, 1211-1218, doi:10.1038/nn.3469 (2013).
- 120 Tay, T. L., Sagar, Dautzenberg, J., Grün, D. & Prinz, M. Unique microglia recovery population revealed by single-cell RNAseq following neurodegeneration. *Acta neuropathologica communications* **6**, 87, doi:10.1186/s40478-018-0584-3 (2018).
- 121 McQuade, A. *et al.* Gene expression and functional deficits underlie TREM2-knockout microglia responses in human models of Alzheimer's disease. *Nature communications* **11**, 5370, doi:10.1038/s41467-020-19227-5 (2020).
- 122 Hickman, S., Izzy, S., Sen, P., Morsett, L. & El Khoury, J. Microglia in neurodegeneration. *Nature neuroscience* **21**, 1359-1369, doi:10.1038/s41593-018-0242-x (2018).
- 123 Cignarella, F. *et al.* TREM2 activation on microglia promotes myelin debris clearance and remyelination in a model of multiple sclerosis. *Acta Neuropathologica* **140**, 513-534, doi:10.1007/s00401-020-02193-z (2020).

- 124 Piccio, L. *et al.* Blockade of TREM-2 exacerbates experimental autoimmune encephalomyelitis. *European journal of immunology* **37**, 1290-1301, doi:10.1002/eji.200636837 (2007).
- 125 Takahashi, K., Prinz, M., Stagi, M., Chechneva, O. & Neumann, H. TREM2-transduced myeloid precursors mediate nervous tissue debris clearance and facilitate recovery in an animal model of multiple sclerosis. *PLoS medicine* **4**, e124, doi:10.1371/journal.pmed.0040124 (2007).
- 126 Konishi, H. & Kiyama, H. Microglial TREM2/DAP12 Signaling: A Double-Edged Sword in Neural Diseases. *Frontiers in cellular neuroscience* **12**, 206, doi:10.3389/fncel.2018.00206 (2018).
- 127 Satoh, J. *et al.* LC3, an autophagosome marker, is expressed on oligodendrocytes in Nasu-Hakola disease brains. *Orphanet journal of rare diseases* **9**, 68, doi:10.1186/1750-1172-9-68 (2014).
- 128 Deczkowska, A., Weiner, A. & Amit, I. The Physiology, Pathology, and Potential Therapeutic Applications of the TREM2 Signaling Pathway. *Cell* **181**, 1207-1217, doi:10.1016/j.cell.2020.05.003 (2020).
- 129 Kober, D. L. & Brett, T. J. TREM2-Ligand Interactions in Health and Disease. *J Mol Biol* **429**, 1607-1629, doi:10.1016/j.jmb.2017.04.004 (2017).
- 130 Peng, Q. *et al.* TREM2- and DAP12-dependent activation of PI3K requires DAP10 and is inhibited by SHIP1. *Science signaling* **3**, ra38, doi:10.1126/scisignal.2000500 (2010).
- 131 Nugent, A. A. *et al.* TREM2 Regulates Microglial Cholesterol Metabolism upon Chronic Phagocytic Challenge. *Neuron* **105**, 837-854.e839, doi:10.1016/j.neuron.2019.12.007 (2020).
- 132 Healy, L. M. *et al.* MerTK-mediated regulation of myelin phagocytosis by macrophages generated from patients with MS. *Neurology(R) neuroimmunology & neuroinflammation* **4**, e402, doi:10.1212/nxi.0000000000000402 (2017).
- 133 Bellan, M., Pirisi, M. & Sainaghi, P. P. The Gas6/TAM System and Multiple Sclerosis. *International journal of molecular sciences* **17**, doi:10.3390/ijms17111807 (2016).
- 134 Grajchen, E. *et al.* CD36-mediated uptake of myelin debris by macrophages and microglia reduces neuroinflammation. *Journal of Neuroinflammation* **17**, 224, doi:10.1186/s12974-020-01899-x (2020).
- 135 Reichert, F. & Rotshenker, S. Complement-receptor-3 and scavenger-receptor-AI/II mediated myelin phagocytosis in microglia and macrophages. *Neurobiology of disease* **12**, 65-72, doi:10.1016/s0969-9961(02)00008-6 (2003).
- 136 Boven, L. A. *et al.* Myelin-laden macrophages are anti-inflammatory, consistent with foam cells in multiple sclerosis. *Brain : a journal of neurology* **129**, 517-526, doi:10.1093/brain/awh707 (2006).
- 137 Li, W. W., Setzu, A., Zhao, C. & Franklin, R. J. Minocycline-mediated inhibition of microglia activation impairs oligodendrocyte progenitor cell responses and remyelination in a non-immune model of demyelination. *Journal of neuroimmunology* **158**, 58-66, doi:10.1016/j.jneuroim.2004.08.011 (2005).
- 138 Laffitte, B. A. *et al.* LXRs control lipid-inducible expression of the apolipoprotein E gene in macrophages and adipocytes. *Proceedings of the National Academy of Sciences of the United States of America* **98**, 507-512, doi:10.1073/pnas.021488798 (2001).
- 139 Savage, J. C. *et al.* Nuclear receptors license phagocytosis by trem2+ myeloid cells in mouse models of Alzheimer's disease. *The Journal of neuroscience : the official*

- Journal of the Society for Neuroscience* **35**, 6532-6543, doi:10.1523/jneurosci.4586-14.2015 (2015).
- 140 Muth, C., Hartmann, A., Sepulveda-Falla, D., Glatzel, M. & Krasemann, S. Phagocytosis of Apoptotic Cells Is Specifically Upregulated in ApoE4 Expressing Microglia in vitro. *Frontiers in cellular neuroscience* **13**, doi:10.3389/fncel.2019.00181 (2019).
- 141 Dean, D. C., 3rd *et al.* Brain differences in infants at differential genetic risk for late-onset Alzheimer disease: a cross-sectional imaging study. *JAMA neurology* **71**, 11-22, doi:10.1001/jamaneurol.2013.4544 (2014).
- 142 Operto, G. *et al.* Interactive effect of age and APOE- ϵ 4 allele load on white matter myelin content in cognitively normal middle-aged subjects. *NeuroImage: Clinical* **24**, 101983, doi:https://doi.org/10.1016/j.nicl.2019.101983 (2019).
- 143 Fazekas, F. *et al.* Apolipoprotein E ϵ 4 is associated with rapid progression of multiple sclerosis. *Neurology* **57**, 853-857, doi:10.1212/wnl.57.5.853 (2001).
- 144 Lill, C. M. *et al.* Closing the case of APOE in multiple sclerosis: no association with disease risk in over 29 000 subjects. *Journal of medical genetics* **49**, 558-562, doi:10.1136/jmedgenet-2012-101175 (2012).
- 145 Schmidt, S. *et al.* Association of Polymorphisms in the Apolipoprotein E Region with Susceptibility to and Progression of Multiple Sclerosis. *The American Journal of Human Genetics* **70**, 708-717, doi:https://doi.org/10.1086/339269 (2002).
- 146 Hagemeyer, N. *et al.* Microglia contribute to normal myelinogenesis and to oligodendrocyte progenitor maintenance during adulthood. *Acta Neuropathol* **134**, 441-458, doi:10.1007/s00401-017-1747-1 (2017).
- 147 Kracht, L. *et al.* Human fetal microglia acquire homeostatic immune-sensing properties early in development. *Science (New York, N.Y.)* **369**, 530-537, doi:10.1126/science.aba5906 (2020).
- 148 Bode, K. *et al.* Dectin-1 Binding to Annexins on Apoptotic Cells Induces Peripheral Immune Tolerance via NADPH Oxidase-2. *Cell reports* **29**, 4435-4446.e4439, doi:10.1016/j.celrep.2019.11.086 (2019).
- 149 Li, H. *et al.* Low dose zymosan ameliorates both chronic and relapsing experimental autoimmune encephalomyelitis. *Journal of neuroimmunology* **254**, 28-38, doi:10.1016/j.jneuroim.2012.08.013 (2013).
- 150 Deerhake, M. E. *et al.* Dectin-1 limits central nervous system autoimmunity through a non-canonical pathway. *bioRxiv*, 2020.2005.2006.080481, doi:10.1101/2020.05.06.080481 (2020).
- 151 Baldwin, K. T., Carbajal, K. S., Segal, B. M. & Giger, R. J. Neuroinflammation triggered by β -glucan/dectin-1 signaling enables CNS axon regeneration. *Proceedings of the National Academy of Sciences of the United States of America* **112**, 2581-2586, doi:10.1073/pnas.1423221112 (2015).
- 152 Diaz-Aparicio, I. *et al.* Microglia Actively Remodel Adult Hippocampal Neurogenesis through the Phagocytosis Secretome. *The Journal of Neuroscience* **40**, 1453, doi:10.1523/JNEUROSCI.0993-19.2019 (2020).
- 153 Sierra, A. *et al.* Microglia shape adult hippocampal neurogenesis through apoptosis-coupled phagocytosis. *Cell stem cell* **7**, 483-495, doi:10.1016/j.stem.2010.08.014 (2010).
- 154 Paolicelli, R. C. *et al.* Synaptic pruning by microglia is necessary for normal brain development. *Science (New York, N.Y.)* **333**, 1456-1458, doi:10.1126/science.1202529 (2011).

- 155 Hughes, A. N. & Appel, B. Microglia phagocytose myelin sheaths to modify developmental myelination. *Nature neuroscience* **23**, 1055-1066, doi:10.1038/s41593-020-0654-2 (2020).
- 156 Pagani, F. *et al.* Defective microglial development in the hippocampus of Cx3cr1 deficient mice. *Frontiers in cellular neuroscience* **9**, 111, doi:10.3389/fncel.2015.00111 (2015).
- 157 Schafer, D. P. *et al.* Microglia sculpt postnatal neural circuits in an activity and complement-dependent manner. *Neuron* **74**, 691-705, doi:10.1016/j.neuron.2012.03.026 (2012).
- 158 Filipello, F. *et al.* The Microglial Innate Immune Receptor TREM2 Is Required for Synapse Elimination and Normal Brain Connectivity. *Immunity* **48**, 979-991.e978, doi:10.1016/j.immuni.2018.04.016 (2018).
- 159 Shi, Q. *et al.* Complement C3-Deficient Mice Fail to Display Age-Related Hippocampal Decline. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **35**, 13029-13042, doi:10.1523/jneurosci.1698-15.2015 (2015).
- 160 Stephan, A. H. *et al.* A dramatic increase of C1q protein in the CNS during normal aging. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **33**, 13460-13474, doi:10.1523/jneurosci.1333-13.2013 (2013).
- 161 Koning, N., Swaab, D. F., Hoek, R. M. & Huitinga, I. Distribution of the immune inhibitory molecules CD200 and CD200R in the normal central nervous system and multiple sclerosis lesions suggests neuron-glia and glia-glia interactions. *J Neuropathol Exp Neurol* **68**, 159-167, doi:10.1097/NEN.0b013e3181964113 (2009).
- 162 Szepesi, Z., Manouchehrian, O., Bachiller, S. & Deierborg, T. Bidirectional Microglia-Neuron Communication in Health and Disease. *Frontiers in cellular neuroscience* **12**, 323-323, doi:10.3389/fncel.2018.00323 (2018).
- 163 Ridderstad Wollberg, A. *et al.* Pharmacological inhibition of the chemokine receptor CX3CR1 attenuates disease in a chronic-relapsing rat model for multiple sclerosis. *Proceedings of the National Academy of Sciences* **111**, 5409-5414, doi:10.1073/pnas.1316510111 (2014).
- 164 Garcia, J. A. *et al.* Regulation of adaptive immunity by the fractalkine receptor during autoimmune inflammation. *Journal of immunology (Baltimore, Md. : 1950)* **191**, 1063-1072, doi:10.4049/jimmunol.1300040 (2013).
- 165 Liu, X., Yao, D. L. & Webster, H. Insulin-like growth factor I treatment reduces clinical deficits and lesion severity in acute demyelinating experimental autoimmune encephalomyelitis. *Multiple sclerosis (Houndmills, Basingstoke, England)* **1**, 2-9, doi:10.1177/135245859500100102 (1995).
- 166 Labandeira-Garcia, J. L., Costa-Besada, M. A., Labandeira, C. M., Villar-Cheda, B. & Rodríguez-Perez, A. I. Insulin-Like Growth Factor-1 and Neuroinflammation. *Front Aging Neurosci* **9**, 365-365, doi:10.3389/fnagi.2017.00365 (2017).
- 167 Chung, E. Y. *et al.* Interleukin-10 Expression in Macrophages during Phagocytosis of Apoptotic Cells Is Mediated by Homeodomain Proteins Pbx1 and Prep-1. *Immunity* **27**, 952-964, doi:https://doi.org/10.1016/j.immuni.2007.11.014 (2007).
- 168 Aarum, J., Sandberg, K., Haerberlein, S. L. B. & Persson, M. A. A. Migration and differentiation of neural precursor cells can be directed by microglia. *Proceedings of the National Academy of Sciences* **100**, 15983-15988, doi:10.1073/pnas.2237050100 (2003).

- 169 Gray, E., Thomas, T. L., Betmouni, S., Scolding, N. & Love, S. Elevated myeloperoxidase activity in white matter in multiple sclerosis. *Neuroscience letters* **444**, 195-198, doi:https://doi.org/10.1016/j.neulet.2008.08.035 (2008).
- 170 Mao, P. & Reddy, P. H. Is multiple sclerosis a mitochondrial disease? *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* **1802**, 66-79, doi:https://doi.org/10.1016/j.bbadis.2009.07.002 (2010).
- 171 Ohl, K., Tenbrock, K. & Kipp, M. Oxidative stress in multiple sclerosis: Central and peripheral mode of action. *Experimental neurology* **277**, 58-67, doi:10.1016/j.expneurol.2015.11.010 (2016).
- 172 Linker, R. A. *et al.* Fumaric acid esters exert neuroprotective effects in neuroinflammation via activation of the Nrf2 antioxidant pathway. *Brain : a journal of neurology* **134**, 678-692, doi:10.1093/brain/awq386 (2011).
- 173 Forte, M. *et al.* Cyclophilin D inactivation protects axons in experimental autoimmune encephalomyelitis, an animal model of multiple sclerosis. *Proceedings of the National Academy of Sciences of the United States of America* **104**, 7558-7563, doi:10.1073/pnas.0702228104 (2007).
- 174 Schluesener, H. J. & Seid, K. Heme oxygenase-1 in lesions of rat experimental autoimmune encephalomyelitis and neuritis. *Journal of neuroimmunology* **110**, 114-120, doi:10.1016/s0165-5728(00)00352-0 (2000).
- 175 Kobayashi, E. H. *et al.* Nrf2 suppresses macrophage inflammatory response by blocking proinflammatory cytokine transcription. *Nature communications* **7**, 11624, doi:10.1038/ncomms11624 (2016).
- 176 Carlström, K. E. *et al.* Gsta4 controls apoptosis of differentiating adult oligodendrocytes during homeostasis and remyelination via the mitochondria-associated Fas-Casp8-Bid-axis. *Nature communications* **11**, 4071, doi:10.1038/s41467-020-17871-5 (2020).
- 177 Choi, B. Y. *et al.* Inhibition of NADPH oxidase activation reduces EAE-induced white matter damage in mice. *J Neuroinflammation* **12**, 104, doi:10.1186/s12974-015-0325-5 (2015).
- 178 Witherick, J., Wilkins, A., Scolding, N. & Kemp, K. Mechanisms of Oxidative Damage in Multiple Sclerosis and a Cell Therapy Approach to Treatment. *Autoimmune Diseases* **2011**, 164608, doi:10.4061/2011/164608 (2011).
- 179 van der Goes, A. *et al.* Reactive oxygen species are required for the phagocytosis of myelin by macrophages. *Journal of neuroimmunology* **92**, 67-75, doi:10.1016/s0165-5728(98)00175-1 (1998).
- 180 Williams, K., Ulvestad, E., Waage, A., Antel, J. P. & McLaurin, J. Activation of adult human derived microglia by myelin phagocytosis in vitro. *Journal of neuroscience research* **38**, 433-443, doi:10.1002/jnr.490380409 (1994).
- 181 Carlström, K. E. *et al.* Therapeutic efficacy of dimethyl fumarate in relapsing-remitting multiple sclerosis associates with ROS pathway in monocytes. *Nature communications* **10**, 3081, doi:10.1038/s41467-019-11139-3 (2019).
- 182 Kang, Z. *et al.* Act1 mediates IL-17-induced EAE pathogenesis selectively in NG2+ glial cells. *Nature neuroscience* **16**, 1401-1408, doi:10.1038/nn.3505 (2013).
- 183 Bonora, M. *et al.* Tumor necrosis factor- α impairs oligodendroglial differentiation through a mitochondria-dependent process. *Cell Death & Differentiation* **21**, 1198-1208, doi:10.1038/cdd.2014.35 (2014).
- 184 Lin, W. *et al.* Interferon-gamma inhibits central nervous system remyelination through a process modulated by endoplasmic reticulum stress. *Brain : a journal of neurology* **129**, 1306-1318, doi:10.1093/brain/awl044 (2006).

- 185 Malhotra, S. *et al.* NLRP3 inflammasome as prognostic factor and therapeutic target in primary progressive multiple sclerosis patients. *Brain : a journal of neurology* **143**, 1414-1430, doi:10.1093/brain/awaa084 (2020).
- 186 Komuczki, J. *et al.* Fate-Mapping of GM-CSF Expression Identifies a Discrete Subset of Inflammation-Driving T Helper Cells Regulated by Cytokines IL-23 and IL-1 β . *Immunity* **50**, 1289-1304.e1286, doi:10.1016/j.immuni.2019.04.006 (2019).
- 187 Hauptmann, J. *et al.* Interleukin-1 promotes autoimmune neuroinflammation by suppressing endothelial heme oxygenase-1 at the blood–brain barrier. *Acta Neuropathologica* **140**, 549-567, doi:10.1007/s00401-020-02187-x (2020).
- 188 Mason, J. L., Suzuki, K., Chaplin, D. D. & Matsushima, G. K. Interleukin-1 β promotes repair of the CNS. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **21**, 7046-7052, doi:10.1523/jneurosci.21-18-07046.2001 (2001).
- 189 Vela, J. M., Molina-Holgado, E., Arévalo-Martín, A., Almazán, G. & Guaza, C. Interleukin-1 regulates proliferation and differentiation of oligodendrocyte progenitor cells. *Molecular and cellular neurosciences* **20**, 489-502, doi:10.1006/mcne.2002.1127 (2002).
- 190 Hauser, S. L., Doolittle, T. H., Lincoln, R., Brown, R. H. & Dinarello, C. A. Cytokine accumulations in CSF of multiple sclerosis patients: frequent detection of interleukin-1 and tumor necrosis factor but not interleukin-6. *Neurology* **40**, 1735-1739, doi:10.1212/wnl.40.11.1735 (1990).
- 191 Titus, H. E. *et al.* Pre-clinical and Clinical Implications of “Inside-Out” vs. “Outside-In” Paradigms in Multiple Sclerosis Etiopathogenesis. *Frontiers in cellular neuroscience* **14**, 599717, doi:10.3389/fncel.2020.599717 (2020).
- 192 Titus, H. E. *et al.* Pre-clinical and Clinical Implications of "Inside-Out" vs. "Outside-In" Paradigms in Multiple Sclerosis Etiopathogenesis. *Frontiers in cellular neuroscience* **14**, 599717-599717, doi:10.3389/fncel.2020.599717 (2020).
- 193 Cloake, N. C., Yan, J., Aminian, A., Pender, M. P. & Greer, J. M. PLP1 Mutations in Patients with Multiple Sclerosis: Identification of a New Mutation and Potential Pathogenicity of the Mutations. *Journal of clinical medicine* **7**, doi:10.3390/jcm7100342 (2018).
- 194 Angelova, D. M. & Brown, D. R. Microglia and the aging brain: are senescent microglia the key to neurodegeneration? *Journal of Neurochemistry* **151**, 676-688, doi:https://doi.org/10.1111/jnc.14860 (2019).
- 195 Shaw, A. C., Goldstein, D. R. & Montgomery, R. R. Age-dependent dysregulation of innate immunity. *Nat. Rev. Immunol.* **13**, 875-887, doi:10.1038/nri3547 (2013).
- 196 Stranks, A. J. *et al.* Autophagy Controls Acquisition of Aging Features in Macrophages. *Journal of innate immunity* **7**, 375-391, doi:10.1159/000370112 (2015).
- 197 Galatro, T. F. *et al.* Transcriptomic analysis of purified human cortical microglia reveals age-associated changes. *Nature neuroscience* **20**, 1162-1171, doi:10.1038/nn.4597 (2017).
- 198 Olah, M. *et al.* A transcriptomic atlas of aged human microglia. *Nature communications* **9**, 539, doi:10.1038/s41467-018-02926-5 (2018).
- 199 Njie, E. G. *et al.* Ex vivo cultures of microglia from young and aged rodent brain reveal age-related changes in microglial function. *Neurobiology of aging* **33**, 195.e191-112, doi:10.1016/j.neurobiolaging.2010.05.008 (2012).

- 200 Safaiyan, S. *et al.* Age-related myelin degradation burdens the clearance function of microglia during aging. *Nature neuroscience* **19**, 995-998, doi:10.1038/nn.4325 (2016).
- 201 Wang, C., Telpoukhovskaia, M. A., Bahr, B. A., Chen, X. & Gan, L. Endo-lysosomal dysfunction: a converging mechanism in neurodegenerative diseases. *Current opinion in neurobiology* **48**, 52-58, doi:10.1016/j.conb.2017.09.005 (2018).
- 202 Hanslik, K. L. & Ulland, T. K. The Role of Microglia and the Nlrp3 Inflammasome in Alzheimer's Disease. *Frontiers in Neurology* **11**, doi:10.3389/fneur.2020.570711 (2020).
- 203 Koellhoffer, E. C., McCullough, L. D. & Ritzel, R. M. Old Maids: Aging and Its Impact on Microglia Function. *International journal of molecular sciences* **18**, doi:10.3390/ijms18040769 (2017).
- 204 Höhn, A. & Grune, T. Lipofuscin: formation, effects and role of macroautophagy. *Redox Biology* **1**, 140-144, doi:https://doi.org/10.1016/j.redox.2013.01.006 (2013).
- 205 Sierra, A., Gottfried-Blackmore, A. C., McEwen, B. S. & Bulloch, K. Microglia derived from aging mice exhibit an altered inflammatory profile. *Glia* **55**, 412-424, doi:10.1002/glia.20468 (2007).
- 206 Elmore, M. R., Lee, R. J., West, B. L. & Green, K. N. Characterizing newly repopulated microglia in the adult mouse: impacts on animal behavior, cell morphology, and neuroinflammation. *PloS one* **10**, e0122912, doi:10.1371/journal.pone.0122912 (2015).
- 207 Willis, E. F. *et al.* Repopulating Microglia Promote Brain Repair in an IL-6-Dependent Manner. *Cell* **180**, 833-846.e816, doi:10.1016/j.cell.2020.02.013 (2020).
- 208 Elmore, M. R. P. *et al.* Replacement of microglia in the aged brain reverses cognitive, synaptic, and neuronal deficits in mice. *Aging Cell* **17**, e12832, doi:https://doi.org/10.1111/acer.12832 (2018).
- 209 De Biase, L. M. *et al.* Local Cues Establish and Maintain Region-Specific Phenotypes of Basal Ganglia Microglia. *Neuron* **95**, 341-356.e346, doi:10.1016/j.neuron.2017.06.020 (2017).
- 210 Utz, S. G. *et al.* Early Fate Defines Microglia and Non-parenchymal Brain Macrophage Development. *Cell* **181**, 557-573.e518, doi:10.1016/j.cell.2020.03.021 (2020).
- 211 Van Hove, H. *et al.* A single-cell atlas of mouse brain macrophages reveals unique transcriptional identities shaped by ontogeny and tissue environment. *Nature neuroscience* **22**, 1021-1035, doi:10.1038/s41593-019-0393-4 (2019).
- 212 Ling, E. A., Kaur, C. & Lu, J. Origin, nature, and some functional considerations of intraventricular macrophages, with special reference to the ependymal cells. *Microscopy research and technique* **41**, 43-56, doi:10.1002/(sici)1097-0029(19980401)41:1<43::Aid-jemt5>3.0.Co;2-v (1998).
- 213 Kierdorf, K., Masuda, T., Jordão, M. J. C. & Prinz, M. Macrophages at CNS interfaces: ontogeny and function in health and disease. *Nature Reviews Neuroscience* **20**, 547-562, doi:10.1038/s41583-019-0201-x (2019).
- 214 Rodríguez-Lorenzo, S. *et al.* Inflammation of the choroid plexus in progressive multiple sclerosis: accumulation of granulocytes and T cells. *Acta neuropathologica communications* **8**, 9, doi:10.1186/s40478-020-0885-1 (2020).
- 215 Magliozzi, R. *et al.* Meningeal B-cell follicles in secondary progressive multiple sclerosis associate with early onset of disease and severe cortical pathology. *Brain : a journal of neurology* **130**, 1089-1104, doi:10.1093/brain/awm038 (2007).

- 216 Cueto, F. J., del Fresno, C. & Sancho, D. DNGR-1, a Dendritic Cell-Specific Sensor of
Tissue Damage That Dually Modulates Immunity and Inflammation. *Frontiers in*
immunology **10**, doi:10.3389/fimmu.2019.03146 (2020).
- 217 Ajami, B. *et al.* Single-cell mass cytometry reveals distinct populations of brain
myeloid cells in mouse neuroinflammation and neurodegeneration models. *Nature*
neuroscience **21**, 541-551, doi:10.1038/s41593-018-0100-x (2018).
- 218 Mildner, A. *et al.* CCR2+Ly-6Chi monocytes are crucial for the effector phase of
autoimmunity in the central nervous system. *Brain : a journal of neurology* **132**,
2487-2500, doi:10.1093/brain/awp144 (2009).
- 219 Fife, B. T., Huffnagle, G. B., Kuziel, W. A. & Karpus, W. J. CC chemokine receptor 2 is
critical for induction of experimental autoimmune encephalomyelitis. *The Journal*
of experimental medicine **192**, 899-905, doi:10.1084/jem.192.6.899 (2000).
- 220 Ajami, B., Bennett, J. L., Krieger, C., McNagny, K. M. & Rossi, F. M. Infiltrating
monocytes trigger EAE progression, but do not contribute to the resident microglia
pool. *Nature neuroscience* **14**, 1142-1149, doi:10.1038/nn.2887 (2011).
- 221 Cao, Y. *et al.* Functional inflammatory profiles distinguish myelin-reactive T cells
from patients with multiple sclerosis. *Science translational medicine* **7**, 287ra274,
doi:10.1126/scitranslmed.aaa8038 (2015).
- 222 Lotfi, N., Zhang, G.-X., Esmaeil, N. & Rostami, A. Evaluation of the effect of GM-CSF
blocking on the phenotype and function of human monocytes. *Scientific reports* **10**,
1567, doi:10.1038/s41598-020-58131-2 (2020).
- 223 Zhang, X. M., Lund, H., Mia, S., Parsa, R. & Harris, R. A. Adoptive transfer of
cytokine-induced immunomodulatory adult microglia attenuates experimental
autoimmune encephalomyelitis in DBA/1 mice. *Glia* **62**, 804-817,
doi:10.1002/glia.22643 (2014).
- 224 Wolf, Y. *et al.* Microglial MHC class II is dispensable for experimental autoimmune
encephalomyelitis and cuprizone-induced demyelination. *European journal of*
immunology **48**, 1308-1318, doi:10.1002/eji.201847540 (2018).
- 225 Wlodarczyk, A., Løbner, M., Cédile, O. & Owens, T. Comparison of microglia and
infiltrating CD11c+ cells as antigen presenting cells for T cell proliferation and
cytokine response. *Journal of Neuroinflammation* **11**, 57, doi:10.1186/1742-2094-
11-57 (2014).
- 226 Mundt, S. *et al.* Conventional DCs sample and present myelin antigens in the
healthy CNS and allow parenchymal T cell entry to initiate neuroinflammation.
Science immunology **4**, doi:10.1126/sciimmunol.aau8380 (2019).
- 227 Monaghan, K. L., Zheng, W., Hu, G. & Wan, E. C. K. Monocytes and Monocyte-
Derived Antigen-Presenting Cells Have Distinct Gene Signatures in Experimental
Model of Multiple Sclerosis. *Frontiers in immunology* **10**,
doi:10.3389/fimmu.2019.02779 (2019).
- 228 Keller, C. W. *et al.* CYBB/NOX2 in conventional DCs controls T cell
encephalitogenicity during neuroinflammation. *Autophagy*, 1-15,
doi:10.1080/15548627.2020.1756678 (2020).
- 229 Ulland, T. K. *et al.* TREM2 Maintains Microglial Metabolic Fitness in Alzheimer's
Disease. *Cell* **170**, 649-663.e613, doi:10.1016/j.cell.2017.07.023 (2017).
- 230 Obba, S. *et al.* The PRKAA1/AMPKα1 pathway triggers autophagy during CSF1-
induced human monocyte differentiation and is a potential target in CMML.
Autophagy **11**, 1114-1129, doi:10.1080/15548627.2015.1034406 (2015).

- 231 Tam, J. M. *et al.* Dectin-1-dependent LC3 recruitment to phagosomes enhances fungicidal activity in macrophages. *The Journal of infectious diseases* **210**, 1844-1854, doi:10.1093/infdis/jiu290 (2014).
- 232 Galloway, D. A., Phillips, A. E. M., Owen, D. R. J. & Moore, C. S. Phagocytosis in the Brain: Homeostasis and Disease. *Frontiers in immunology* **10**, doi:10.3389/fimmu.2019.00790 (2019).
- 233 Kang, R., Zeh, H. J., Lotze, M. T. & Tang, D. The Beclin 1 network regulates autophagy and apoptosis. *Cell death and differentiation* **18**, 571-580, doi:10.1038/cdd.2010.191 (2011).
- 234 Keane, L. *et al.* mTOR-dependent translation amplifies microglia priming in aging mice. *The Journal of clinical investigation*, doi:10.1172/jci132727 (2020).
- 235 Zachari, M. & Ganley, I. G. The mammalian ULK1 complex and autophagy initiation. *Essays in biochemistry* **61**, 585-596, doi:10.1042/ebc20170021 (2017).
- 236 Wirth, M. *et al.* Molecular determinants regulating selective binding of autophagy adapters and receptors to ATG8 proteins. *Nature communications* **10**, 2055, doi:10.1038/s41467-019-10059-6 (2019).
- 237 Lőrincz, P. & Juhász, G. Autophagosome-Lysosome Fusion. *Journal of Molecular Biology* **432**, 2462-2482, doi:https://doi.org/10.1016/j.jmb.2019.10.028 (2020).
- 238 Kuma, A., Komatsu, M. & Mizushima, N. Autophagy-monitoring and autophagy-deficient mice. *Autophagy* **13**, 1619-1628, doi:10.1080/15548627.2017.1343770 (2017).
- 239 Mortensen, M. *et al.* The autophagy protein Atg7 is essential for hematopoietic stem cell maintenance. *The Journal of experimental medicine* **208**, 455-467, doi:10.1084/jem.20101145 (2011).
- 240 Clarke, A. J. & Simon, A. K. Autophagy in the renewal, differentiation and homeostasis of immune cells. *Nat. Rev. Immunol.* **19**, 170-183, doi:10.1038/s41577-018-0095-2 (2019).
- 241 Keller, C. W. *et al.* ATG-dependent phagocytosis in dendritic cells drives myelin-specific CD4(+) T cell pathogenicity during CNS inflammation. *Proceedings of the National Academy of Sciences of the United States of America* **114**, E11228-e11237, doi:10.1073/pnas.1713664114 (2017).
- 242 Levine, B., Mizushima, N. & Virgin, H. W. Autophagy in immunity and inflammation. *Nature* **469**, 323-335, doi:10.1038/nature09782 (2011).
- 243 Nie, T. *et al.* Regulation of ER stress-induced autophagy by GSK3 β -TIP60-ULK1 pathway. *Cell death & disease* **7**, e2563, doi:10.1038/cddis.2016.423 (2016).
- 244 Rashid, H. O., Yadav, R. K., Kim, H. R. & Chae, H. J. ER stress: Autophagy induction, inhibition and selection. *Autophagy* **11**, 1956-1977, doi:10.1080/15548627.2015.1091141 (2015).
- 245 Komatsu, M. *et al.* Impairment of starvation-induced and constitutive autophagy in Atg7-deficient mice. *The Journal of cell biology* **169**, 425-434, doi:10.1083/jcb.200412022 (2005).
- 246 Kundu, M. *et al.* Ulk1 plays a critical role in the autophagic clearance of mitochondria and ribosomes during reticulocyte maturation. *Blood* **112**, 1493-1502, doi:10.1182/blood-2008-02-137398 (2008).
- 247 Zhang, J. *et al.* Mitochondrial clearance is regulated by Atg7-dependent and -independent mechanisms during reticulocyte maturation. *Blood* **114**, 157-164, doi:10.1182/blood-2008-04-151639 (2009).
- 248 Hametner, S. *et al.* Iron and neurodegeneration in the multiple sclerosis brain. *Annals of neurology* **74**, 848-861, doi:https://doi.org/10.1002/ana.23974 (2013).

- 249 Sanjuan, M. A. *et al.* Toll-like receptor signalling in macrophages links the autophagy pathway to phagocytosis. *Nature* **450**, 1253-1257, doi:10.1038/nature06421 (2007).
- 250 Wong, S. W., Sil, P. & Martinez, J. Rubicon: LC3-associated phagocytosis and beyond. *Febs j* **285**, 1379-1388, doi:10.1111/febs.14354 (2018).
- 251 Ma, J., Becker, C., Lowell, C. A. & Underhill, D. M. Dectin-1-triggered recruitment of light chain 3 protein to phagosomes facilitates major histocompatibility complex class II presentation of fungal-derived antigens. *The Journal of biological chemistry* **287**, 34149-34156, doi:10.1074/jbc.M112.382812 (2012).
- 252 Kyrmizi, I. *et al.* Corticosteroids block autophagy protein recruitment in *Aspergillus fumigatus* phagosomes via targeting dectin-1/Syk kinase signaling. *Journal of immunology (Baltimore, Md. : 1950)* **191**, 1287-1299, doi:10.4049/jimmunol.1300132 (2013).
- 253 Martinez, J. *et al.* Molecular characterization of LC3-associated phagocytosis reveals distinct roles for Rubicon, NOX2 and autophagy proteins. *Nature cell biology* **17**, 893-906, doi:10.1038/ncb3192 (2015).
- 254 Martinez, J. *et al.* Microtubule-associated protein 1 light chain 3 alpha (LC3)-associated phagocytosis is required for the efficient clearance of dead cells. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 17396-17401, doi:10.1073/pnas.1113421108 (2011).
- 255 Heckmann, B. L. *et al.* LC3-Associated Endocytosis Facilitates beta-Amyloid Clearance and Mitigates Neurodegeneration in Murine Alzheimer's Disease. *Cell* **178**, 536-551.e514, doi:10.1016/j.cell.2019.05.056 (2019).
- 256 Lucin, K. M. *et al.* Microglial beclin 1 regulates retromer trafficking and phagocytosis and is impaired in Alzheimer's disease. *Neuron* **79**, 873-886, doi:10.1016/j.neuron.2013.06.046 (2013).
- 257 Martinez, J. *et al.* Noncanonical autophagy inhibits the autoinflammatory, lupus-like response to dying cells. *Nature* **533**, 115-119, doi:10.1038/nature17950 (2016).
- 258 Cheng, J. *et al.* Microglial autophagy defect causes parkinson disease-like symptoms by accelerating inflammasome activation in mice. *Autophagy*, 1-13, doi:10.1080/15548627.2020.1719723 (2020).
- 259 Kim, H. J. *et al.* Deficient autophagy in microglia impairs synaptic pruning and causes social behavioral defects. *Molecular Psychiatry* **22**, 1576-1584, doi:10.1038/mp.2016.103 (2017).
- 260 Choi, I. *et al.* Microglia clear neuron-released α -synuclein via selective autophagy and prevent neurodegeneration. *Nature communications* **11**, 1386, doi:10.1038/s41467-020-15119-w (2020).
- 261 Levine, B. & Kroemer, G. Biological Functions of Autophagy Genes: A Disease Perspective. *Cell* **176**, 11-42, doi:https://doi.org/10.1016/j.cell.2018.09.048 (2019).
- 262 Harris, H. & Rubinsztein, D. C. Control of autophagy as a therapy for neurodegenerative disease. *Nat. Rev. Neurol.* **8**, 108-117, doi:10.1038/nrneurol.2011.200 (2012).
- 263 Nilsson, P. *et al.* A β secretion and plaque formation depend on autophagy. *Cell reports* **5**, 61-69, doi:10.1016/j.celrep.2013.08.042 (2013).
- 264 Hou, X., Watzlawik, J. O., Fiesel, F. C. & Springer, W. Autophagy in Parkinson's Disease. *Journal of Molecular Biology* **432**, 2651-2672, doi:https://doi.org/10.1016/j.jmb.2020.01.037 (2020).

- 265 Kosmidou, M. *et al.* Multiple sclerosis and inflammatory bowel diseases: a systematic review and meta-analysis. *Journal of neurology* **264**, 254-259, doi:10.1007/s00415-016-8340-8 (2017).
- 266 Liu, X., Qin, H. & Xu, J. The role of autophagy in the pathogenesis of systemic lupus erythematosus. *International immunopharmacology* **40**, 351-361, doi:10.1016/j.intimp.2016.09.017 (2016).
- 267 Tam, R. C. *et al.* Human CLEC16A regulates autophagy through modulating mTOR activity. *Experimental cell research* **352**, 304-312, doi:10.1016/j.yexcr.2017.02.017 (2017).
- 268 Redmann, V. *et al.* Clec16a is Critical for Autolysosome Function and Purkinje Cell Survival. *Scientific reports* **6**, 23326, doi:10.1038/srep23326 (2016).
- 269 Soleimanpour, S. A. *et al.* The diabetes susceptibility gene Clec16a regulates mitophagy. *Cell* **157**, 1577-1590, doi:10.1016/j.cell.2014.05.016 (2014).
- 270 van Luijn, M. M. *et al.* Multiple sclerosis-associated CLEC16A controls HLA class II expression via late endosome biogenesis. *Brain : a journal of neurology* **138**, 1531-1547, doi:10.1093/brain/awv080 (2015).
- 271 Galluzzi, L., Bravo-San Pedro, J. M., Levine, B., Green, D. R. & Kroemer, G. Pharmacological modulation of autophagy: therapeutic potential and persisting obstacles. *Nature reviews. Drug discovery* **16**, 487-511, doi:10.1038/nrd.2017.22 (2017).
- 272 Chen, M. J. *et al.* Microglial ERK signaling is a critical regulator of pro-inflammatory immune responses in Alzheimer's disease. *bioRxiv*, 798215, doi:10.1101/798215 (2019).
- 273 Li, Z., Nie, L., Chen, L., Sun, Y. & Li, G. Rapamycin relieves inflammation of experimental autoimmune encephalomyelitis by altering the balance of Treg/Th17 in a mouse model. *Neuroscience letters* **705**, 39-45, doi:10.1016/j.neulet.2019.04.035 (2019).
- 274 Madeo, F., Eisenberg, T., Pietrocola, F. & Kroemer, G. Spermidine in health and disease. *Science (New York, N.Y.)* **359**, doi:10.1126/science.aan2788 (2018).
- 275 Eisenberg, T. *et al.* Cardioprotection and lifespan extension by the natural polyamine spermidine. *Nature medicine* **22**, 1428-1438, doi:10.1038/nm.4222 (2016).
- 276 Guo, X. *et al.* Spermidine alleviates severity of murine experimental autoimmune encephalomyelitis. *Investigative ophthalmology & visual science* **52**, 2696-2703, doi:10.1167/iovs.10-6015 (2011).
- 277 Yang, Q. *et al.* Spermidine alleviates experimental autoimmune encephalomyelitis through inducing inhibitory macrophages. *Cell death and differentiation* **23**, 1850-1861, doi:10.1038/cdd.2016.71 (2016).
- 278 Evans, T. D., Jeong, S. J., Zhang, X., Sergin, I. & Razani, B. TFEB and trehalose drive the macrophage autophagy-lysosome system to protect against atherosclerosis. *Autophagy* **14**, 724-726, doi:10.1080/15548627.2018.1434373 (2018).
- 279 Rusmini, P. *et al.* Trehalose induces autophagy via lysosomal-mediated TFEB activation in models of motoneuron degeneration. *Autophagy*, 1-21, doi:10.1080/15548627.2018.1535292 (2018).
- 280 Sergin, I. *et al.* Exploiting macrophage autophagy-lysosomal biogenesis as a therapy for atherosclerosis. *Nature communications* **8**, 15750, doi:10.1038/ncomms15750 (2017).
- 281 Cortes, C. J. & La Spada, A. R. TFEB dysregulation as a driver of autophagy dysfunction in neurodegenerative disease: Molecular mechanisms, cellular

- processes, and emerging therapeutic opportunities. *Neurobiology of disease* **122**, 83-93, doi:10.1016/j.nbd.2018.05.012 (2019).
- 282 Hosseinpour-Moghaddam, K., Caraglia, M. & Sahebkar, A. Autophagy induction by
trehalose: Molecular mechanisms and therapeutic impacts. *Journal of cellular*
283 *physiology* **233**, 6524-6543, doi:10.1002/jcp.26583 (2018).
- 283 DeBosch, B. J. *et al.* Trehalose inhibits solute carrier 2A (SLC2A) proteins to induce
autophagy and prevent hepatic steatosis. *Science signaling* **9**, ra21-ra21,
doi:10.1126/scisignal.aac5472 (2016).
- 284 Portbury, S. D. *et al.* Trehalose Improves Cognition in the Transgenic Tg2576
Mouse Model of Alzheimer's Disease. *J Alzheimers Dis* **60**, 549-560,
doi:10.3233/jad-170322 (2017).
- 285 Beecham, A. H. *et al.* Analysis of immune-related loci identifies 48 new
susceptibility variants for multiple sclerosis. *Nature genetics* **45**, 1353-1360,
doi:10.1038/ng.2770 (2013).
- 286 Orthgiess, J. *et al.* Neurons exhibit Lyz2 promoter activity in vivo: Implications for
using LysM-Cre mice in myeloid cell research. *European journal of immunology* **46**,
1529-1532, doi:10.1002/eji.201546108 (2016).
- 287 Chappell-Maor, L. *et al.* Comparative analysis of CreER transgenic mice for the
study of brain macrophages: A case study. *European journal of immunology* **50**,
353-362, doi:10.1002/eji.201948342 (2020).
- 288 Parkhurst, C. N. *et al.* Microglia promote learning-dependent synapse formation
through brain-derived neurotrophic factor. *Cell* **155**, 1596-1609,
doi:10.1016/j.cell.2013.11.030 (2013).
- 289 Dumas, A. *et al.* The inflammasome pyrin contributes to pertussis toxin-induced IL-
1 β synthesis, neutrophil intravascular crawling and autoimmune
encephalomyelitis. *PLoS pathogens* **10**, e1004150,
doi:10.1371/journal.ppat.1004150 (2014).
- 290 Tigno-Aranjuez, J. T., Jaini, R., Tuohy, V. K., Lehmann, P. V. & Tary-Lehmann, M.
Encephalitogenicity of Complete Freund's Adjuvant Relative to CpG Is Linked to
Induction of Th17 Cells. *The Journal of Immunology* **183**, 5654-5661,
doi:10.4049/jimmunol.0900645 (2009).
- 291 Shenderov, K. *et al.* Inflammasome-dependent IL-1 β production is critical for
complete Freund's adjuvant-induced helper T cell polarization (136.44). *The*
Journal of Immunology **184**, 136.144-136.144 (2010).
- 292 Caldeira, C. *et al.* Microglia change from a reactive to an age-like phenotype with
the time in culture. *Frontiers in cellular neuroscience* **8**, 152,
doi:10.3389/fncel.2014.00152 (2014).
- 293 Demeter, A. *et al.* ULK1 and ULK2 are less redundant than previously thought:
computational analysis uncovers distinct regulation and functions of these
autophagy induction proteins. *Scientific reports* **10**, 10940, doi:10.1038/s41598-
020-67780-2 (2020).
- 294 Thessen Hedreul, M. *et al.* Combining genetic mapping with genome-wide
expression in experimental autoimmune encephalomyelitis highlights a gene
network enriched for T cell functions and candidate genes regulating
autoimmunity. *Hum Mol Genet* **22**, 4952-4966, doi:10.1093/hmg/ddt343 (2013).
- 295 Plaza-Zabala, A., Sierra-Torre, V. & Sierra, A. Autophagy and Microglia: Novel
Partners in Neurodegeneration and Aging. *International journal of molecular*
sciences **18**, doi:10.3390/ijms18030598 (2017).

- 296 Martinez-Lopez, N., Athonvarangkul, D. & Singh, R. Autophagy and aging. *Adv Exp Med Biol* **847**, 73-87, doi:10.1007/978-1-4939-2404-2_3 (2015).
- 297 Martinez-Lopez, N., Athonvarangkul, D., Mishall, P., Sahu, S. & Singh, R. Autophagy proteins regulate ERK phosphorylation. *Nature communications* **4**, 2799, doi:10.1038/ncomms3799 (2013).
- 298 Gao, T. *et al.* Transcriptional regulation of homeostatic and disease-associated-microglial genes by IRF1, LXR β , and CEBP α . *Glia* **67**, 1958-1975, doi:10.1002/glia.23678 (2019).
- 299 Masuda, T. *et al.* Novel Hexb-based tools for studying microglia in the CNS. *Nature immunology* **21**, 802-815, doi:10.1038/s41590-020-0707-4 (2020).
- 300 Gao, H. *et al.* Opposing Functions of Microglial and Macrophagic TNFR2 in the Pathogenesis of Experimental Autoimmune Encephalomyelitis. *Cell reports* **18**, 198-212, doi:10.1016/j.celrep.2016.11.083 (2017).